

Massey products in differential cohomology via stacks

Daniel Grady and Hisham Sati

May 6, 2016

Abstract

We extend Massey products from cohomology to differential cohomology via stacks, organizing and generalizing existing constructions in Deligne cohomology. We study the properties and show how they are related to more classical Massey products in de Rham, singular, and Deligne cohomology. The setting and the algebraic machinery via stacks allow for computations and make the construction well-suited for applications. We illustrate with several examples from differential geometry and mathematical physics.

Contents

1	Introduction	1
2	Massey products and differential cohomology	3
2.1	Classical (generalized) Massey products	3
2.2	Differential cohomology	4
2.3	Cup product in differential cohomology	13
2.4	Massey products in hypercohomology	15
3	Massey products in the language of higher stacks	18
3.1	The Dold-Kan correspondence	18
3.2	Stacky Massey products	20
3.3	General stacky Massey products	24
3.4	Properties of stacky Massey products	27
4	Applications	35
4.1	Trivializations for (higher) structures	35
4.2	Characteristic forms and anomaly cancellations	38
4.3	Transfer of Massey products	41
4.4	Twisted cohomology and twisted Bianchi identities	42
4.5	Quadruple Massey products	44

1 Introduction

Massey products were introduced in [33] and further developed and generalized in [30] [35]. The existence of (higher) Massey products indicates the complexity of the topology of a space. They

also determine whether and how various characterizing properties of a space might be related, in particular how homotopy of a space might be related to its cohomology [28]. On the one hand, Massey products can be viewed as secondary cohomology operations associated with the primary operation given by the cup product. On the other hand, they can also be seen as higher order products in homotopy (A_∞) algebras.

A differential graded algebra (DGA) is a (not a priori commutative) graded algebra A with a map $d : A \rightarrow A$ of degree $+1$ which satisfies the relations (up to sign conventions) $dd = 0$ and $d(ab) = (da)b + (-1)^{\dim a}a(db)$. Then the cohomology $H(A)$ of A with respect to d is a graded algebra. It has further certain operations called (matrix) Massey products, the simplest of which is a correspondence

$$H(A) \otimes H(A) \otimes H(A) \rightarrow H(A) \quad (1.1)$$

which is denoted by $\langle a, b, c \rangle$, where $a, b, c \in H(A)$. This has dimension $\dim(a) + \dim(b) + \dim(c) - 1$, is defined only when $ab = bc = 0 \in H(A)$, and is not well-defined but rather only defined modulo terms of the form $ax + yb$ where x and y are some (auxiliary) elements of $H(A)$. The indeterminacy may, however, sometimes be excluded, for example for dimension reasons, which occurs in applications. Generally, we have $ab = dy$ and $bc = dz$ for $y, z \in A$, so that

$$\langle a, b, c \rangle = yc + (-1)^{\dim a + 1}az \quad (1.2)$$

is a cocycle, with the cohomology class defined modulo the indeterminacy given above.

There are other notions of Massey products, but all are essentially variations on this principle. If the Massey product $\langle a_1, \dots, a_n \rangle$ exists, then all “lower” Massey products necessarily vanish, although the converse is not true in general. One may also apply a similar construction for matrices of elements, leading to matrix Massey products [35], where notions related to formal flatness become important.

Differential cohomology has played an important role recently by combining geometric and topological data, namely usual cohomology and differential forms, in a coherent way [9] [18] [5] [14] [23] [29] [45] [6] [7] [47] [2]. It is natural then to try to extend Massey products, which exist in both of these ingredients, to differential cohomology. Massey products have been considered in Deligne cohomology in [16] [44] [48] [38]. We extend the definitions and constructions to the level of stacks, which has the virtue of allowing for vast generalization to various settings and to a plethora of applications. We believe this formulation has an advantage both for theory and for applications. In particular, we emphasize that desired properties and behaviour of the Massey products are clearer and systematic in stacks, and computations are generally doable and are more efficient there, making them quite suitable for applications. The constructions are based on the thesis [27], but we have taken the opportunity here to sharpen the results and add properties and applications. We emphasize that this paper is the first part of a bigger project, aimed at developing various concrete computational techniques for differential cohomology theories.

The paper is organized as follows. In Sec. 2 we provide the setting for the two main ingredients that we would like to combine, namely classical (generalized) Massey products in Sec. 2.1 and differential cohomology in Sec. 2.2. We set up the former in the general framework of [35] [30] [1] and the latter in the language of stacks (see [21] [43] [47]). Then we recall the Deligne-Beilinson cup product in differential cohomology, as set forth in [19] [20], in Sec. 2.3. The first encounter of Massey products in differential cohomology in the particular setting of Deligne cohomology is

recalled in Sec. 2.4 and adapted slightly to our language. Our main construction is then described in Sec. 3, where we first set up the powerful machinery needed, in the form of the Dold-Kan correspondence, in Sec. 3.1, and then provide the main definitions in Sec. 3.2. Part of this construction, together with a lot of the homotopic background appeared in the second author's thesis [27]. A vast generalization, along the lines of the classic work of May, is presented in Sec. 3.3, where we present all three of differential, singular, and de Rham Massey products within the same setting. Then in Sec. 3.4 we give the properties of the Massey products thus defined. These turn out to be rather attractive in general, with some unexpected features.

In Sec. 4, we illustrate (some aspects of) the construction with various applications. We will first give instances of where the classical Massey products arise in applications, and we apply our constructions in previous sections to supply the differential refinements of these applications. We extend the constructions and discussion in [19] from cup product Chern-Simons theories to what we might call *Massey product Chern-Simons theories*. In Sec. 4.1 we illustrate how (stacky) Massey products arise in trivialization of higher structures, such as (differential) String, Fivebrane [42], and Ninebrane structures [41]. Then in Sec. 4.2 we see how systems arising generally in anomaly cancellation lead naturally to (stacky) Massey products. There are two expressions that involve three differential cohomology classes, namely the stacky Massey product and the triple Deligne-Beilinson cup product. A natural question is whether these are related. Indeed, we establish such a relation via transfer in the context of cobordism in section 4.3. Finally, in settings inspired by type IIA and type IIB string theory in Sec. 4.4 and Sec. 4.5, respectively, we illustrate how these lead to stacky Massey products. Interestingly, the latter gives rise to a quadruple Massey product.

2 Massey products and differential cohomology

2.1 Classical (generalized) Massey products

We recall some notions from [35] [30] [1]. This will be useful for the applications that we will consider later as well as a starting point for comparison with our stacky constructions.

Let (\mathcal{A}, d) be a differential graded algebra over \mathbb{R} endowed with augmentation. Let $M(\mathcal{A})$ be the set of all upper triangular half-infinite matrices with entries in \mathcal{A} , zeroes on the diagonal and finitely many nonzero entries, i.e.

$$M(\mathcal{A}) = \{A = (a_{ij}), a_{ij} \in \mathcal{A}, a_{ij} = 0 \text{ for } j \leq i \text{ and } i, j \geq n + 1 \text{ for some } n\} . \quad (2.1)$$

The last condition distinguishes in $M(\mathcal{A})$ a subset (which is in fact a subalgebra) $M_n(\mathcal{A})$ consisting of all $(n \times n)$ -matrices with entries in \mathcal{A} . The algebra $M(\mathcal{A})$ is bigraded and endowed with a bigraded Lie bracket. We introduce the differential d on $M(\mathcal{A})$ as $dA = (da_{ij})_{i,j \geq 1}$. The algebra \mathcal{A} admits an involution given by $a \mapsto \bar{a} = (-1)^k a$, which can be extended to an automorphism of $M(\mathcal{A})$ as $\bar{A} = (\bar{a}_{ij})_{i,j \geq 1}$, with the differential d satisfying the generalized Leibnitz rule $d(AB) = (dA)B + \bar{A}(dB)$. In [1], the Maurer-Cartan operator $\mu : M \rightarrow M$ was defined as $\mu(A) = dA - \bar{A} \cdot A$. Then a matrix $A \in M$ is said to be a matrix of formal connection if it satisfies the Maurer-Cartan equation in \mathcal{A} ,

$$dA - \bar{A} \cdot A \equiv 0 \text{ mod ker } A , \quad (2.2)$$

i.e. A is a formal connection if $\mu(A) \in \ker A$, where $\ker A$ is a \mathcal{A} -module generated by matrices 1_{ij} such that $A \cdot 1_{ij} = 1_{ij} \cdot A$ (which implies that $AB = BA$ for any matrix $B \in \ker A$). The element

$\mu(A)$ is called the curvature of the formal connection A , and can be shown to be closed (see e.g. [1] [10]).

Now comes the relation between Maurer-Cartan and Massey products. The generalized Massey products are the cohomology classes of the curvature matrices of the formal connection A , i.e. if A is a solution to the Maurer-Cartan equation then the entries of the matrix $[\mu(A)]$ are the generalized Massey products [1]. Geometrically, this means that the latter measure the deviation of connections from flat ones, so that the connection is flat if they vanish. Later we will make use of this approach in describing Massey products in stacks.

Classical Massey products in integral cohomology $H^*(X; \mathbb{Z})$ arise by taking \mathcal{A} to be an algebra over the commutative ring \mathbb{Z} , with the multiplication being associative but not necessarily graded-commutative. Now let α, β, γ be the cohomology classes of closed elements $a \in \mathcal{A}^p$, $b \in \mathcal{A}^q$, and $c \in \mathcal{A}^r$. The triple Massey product $\langle \alpha, \beta, \gamma \rangle$ is defined if one can solve the Maurer-Cartan equation with the formal connection

$$A = \begin{pmatrix} 0 & a & \tilde{f} & * \\ 0 & 0 & b & \tilde{g} \\ 0 & 0 & 0 & c \\ 0 & 0 & 0 & 0 \end{pmatrix}.$$

This is equivalent to the two separate equations

$$d\tilde{f} = (-1)^p a \wedge b \quad \text{and} \quad d\tilde{g} = (-1)^q a \wedge c \quad (2.3)$$

and that implies that the Massey product is defined if and only if

$$\alpha \cup \beta = \beta \cup \gamma = 0 \in H^*(\mathcal{A}). \quad (2.4)$$

The matrix $\mu(A)$ has the form

$$\mu(A) = dA - \overline{A} \cdot A = \begin{pmatrix} 0 & 0 & 0 & \tau \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}$$

and defines the Massey product $[\mu(A)]$ which is equal to the cohomology class

$$\langle \alpha, \beta, \gamma \rangle = [\tau] = \left[(-1)^{p+1} a \wedge \tilde{g} + (-1)^{p+q} \tilde{f} \wedge c \right]. \quad (2.5)$$

Here $[a] \in H^*(\mathcal{A})$ denotes the cohomology class of a closed element $a \in \mathcal{A}$, and $[A] = ([a_{ij}])_{i,j \geq 1} \in M(H^*(\mathcal{A}))$, for a closed matrix $A \in M(\mathcal{A})$, denotes the corresponding matrix whose entries are the cohomology classes of the entries a_{ij} of A . Since \tilde{f} and \tilde{g} are defined by expressions (2.3) up to closed elements from \mathcal{A} , the triple Massey product $\langle \alpha, \beta, \gamma \rangle$ is defined modulo $\alpha \cdot H^{q+r}(\mathcal{A}) + \gamma \cdot H^{p+q}(\mathcal{A})$.

2.2 Differential cohomology

There are several different approaches to differential cohomology. Initially, we will be concerned with the construction as Deligne cohomology [5] [23]. We will then move to the stacky setting,

which illuminates the true nature of differential cohomology as a theory which counts isomorphism classes of higher bundles (and higher abelian gerbes) with connection.

The classical construction relies on hypercohomology of a complex of objects of an abelian category as an extension to complexes of the usual cohomology of an object. For $n \in \mathbb{N}$, let $\mathbb{Z}_{\mathcal{D}}^{\infty}[n]$ be the sheaf of chain complexes given by

$$\mathbb{Z}_{\mathcal{D}}^{\infty}[n] := [\dots \rightarrow 0 \rightarrow \underline{\mathbb{Z}} \hookrightarrow \Omega^0 \rightarrow \Omega^1 \rightarrow \dots \rightarrow \Omega^{n-1}] ,$$

where $\underline{\mathbb{Z}}$ is in degree n and Ω^{n-1} is the sheaf of real-valued $(n-1)$ -forms in degree 0. Given a manifold X , the degree n sheaf hypercohomology with coefficients in $\mathbb{Z}_{\mathcal{D}}^{\infty}[n]$ can be defined to be the degree n differential cohomology of X :

$$\hat{H}^n(X; \mathbb{Z}) := H^n(X; \mathbb{Z}_{\mathcal{D}}^{\infty}[n]) . \quad (2.6)$$

If X is paracompact, then these cohomology groups are given by the cohomology of the total complex of the Čech-Deligne double complex corresponding to a good open cover of X . In what follows, we will always assume that X is paracompact, so that the hypercohomology groups can be computed by either taking arbitrary injective resolutions, or via this more explicit Čech approach.

In [45] (see also [6]), it was observed that these cohomology groups fit nicely into an exact hexagon

$$\begin{array}{ccccc} & \Omega^{n-1}(X)/\text{im}(d) & \xrightarrow{d} & \Omega_{\text{cl}}^n(X) & \\ & \searrow a & & \nearrow R & \\ H_{\text{dR}}^{n-1}(X) & & \hat{H}^n(X; \mathbb{Z}) & & H_{\text{dR}}^n(X) , \\ & \nearrow & \searrow I & & \\ & H^{n-1}(X; U(1)) & \xrightarrow{\quad} & H^n(X, \mathbb{Z}) & \end{array} \quad (2.7)$$

where the bottom row is the Bockstein sequence and the diagonals are exact. The map R is called the *curvature* map and I is called the *integration* map. Notice that, by exactness, in the case that the curvature of a differential cohomology class vanishes, the class lies in the image of the inclusion $H^{n-1}(X; U(1)) \hookrightarrow \hat{H}^n(X; \mathbb{Z})$. We call these classes *flat*, as they represent n -gerbes with connections of vanishing curvature. Differential cohomology therefore detects the topological information when the class is flat and the differential geometric information encoded by the curvature. See [9] [18] [5] [14] [23] [29] [45] [7] [47] [2] for more details on the various approaches.

As we mentioned earlier, our point of view henceforth will be mainly that of stacks. We will recall and introduce some stacks that will be useful for us. We start by surveying some basic concepts and definitions from [32] [21] [19] [20], adapted to our setting. We will provide only as much detail as necessary to introduce our stacks.

Let CartSp be the category with objects copies of Cartesian space \mathbb{R}^n , and morphisms smooth functions. A *smooth prestack* is simply a functor

$$\mathcal{F} : \text{CartSp}^{\text{op}} \rightarrow \text{sSet}$$

with target the category of simplicial sets. The passage from prestacks to stacks is achieved by imposing a sort of gluing condition on \mathcal{F} . Roughly speaking, a stack \mathcal{F} attaches an entire space (equivalently simplicial set) of data to each object in \mathbf{CartSp} . This data should be viewed as being *local* data. The gluing condition then assembles this data into a geometric object, which is a stack. More precisely, we say that a prestack \mathcal{F} satisfies *descent* if for each $U \in \mathbf{CartSp}$ and each open cover $\{U_i\}_{i \in I}$ of U with contractible finite intersections $U_{i_1 i_2 \dots i_k}$, we have a weak equivalence

$$\mathcal{F}(U) \simeq \operatorname{holim} \left\{ \dots \rightrightarrows \prod_{i,j,k} \mathcal{F}(U_{ijk}) \rightrightarrows \prod_{i,j} \mathcal{F}(U_{ij}) \rightrightarrows \prod_i \mathcal{F}(U_i) \right\}. \quad (2.8)$$

In particular, if \mathcal{F} takes values in Kan complexes, this weak equivalence is part of an actual *homotopy* equivalence. The reader may notice the following.

- If we change the target category to \mathbf{Set} and impose the stronger condition that the *strict* limit over the diagram was *isomorphic* to $\mathcal{F}(U)$, we would recover the gluing condition for a sheaf.
- If we change the target category to groupoids, then the above condition recovers the usual notion of descent for classical stacks.

In the latter, homotopy equivalence is simply categorical equivalence of groupoids. Hence the gluing condition respects the correct notion of equivalence (which is weaker than isomorphism). We can therefore view the equivalence (2.8) as the more general gluing condition for ∞ -groupoids (or Kan complexes).

Definition 1. We call a smooth prestack \mathcal{F} a smooth stack if it satisfies descent. We denote the full subcategory of smooth stacks by

$$\mathbf{Sh}_\infty(\mathbf{CartSp}) \hookrightarrow [\mathbf{CartSp}^{\mathrm{op}}, \mathbf{sSet}].$$

Remark 1. The inclusion functor admits a left adjoint L which preserves homotopy colimits (in fact, a left Quillen adjoint [47]). We call this functor L the stackification functor and call the image of a prestack \mathcal{F} under L the stackification of \mathcal{F} .

In [21], the moduli stack of n -gerbes with connection, $\mathbb{B}^n U(1)_{\mathrm{conn}}$, was introduced. This stack was obtained as the stackification of the n -prestack obtained by applying the Dold-Kan map (see section 3.1) to the Deligne presheaf of chain complexes

$$\mathbb{Z}_{\mathcal{D}}^\infty[n+1] := [\dots 0 \rightarrow \mathbb{Z} \hookrightarrow \Omega^1 \rightarrow \Omega^2 \rightarrow \dots \rightarrow \Omega^n].$$

These stacks are the differential analogues of Eilenberg-MacLane spaces and, for a fixed manifold X , there is a bijective correspondence (a ‘representation’)

$$\widehat{H}_{\mathcal{D}}^{n+1}(X; \mathbb{Z}) \simeq \pi_0 \operatorname{Map}(X, \mathbb{B}^n U(1)_{\mathrm{conn}}), \quad (2.9)$$

where the right hand side is the set of morphisms in the homotopy category of stacks.

Remark 2. In general the right hand side of the correspondence (2.9) may not be well-defined. In order to be able to take homotopy groups of the mapping space $Y := \operatorname{Map}(X, \mathbb{B}^n U(1)_{\mathrm{conn}})$, Y has to be a Kan complex, which is the case when X is cofibrant and $\mathbb{B}^n U(1)_{\mathrm{conn}}$ is fibrant. However,

since $\mathbb{B}^n U(1)_{\text{conn}}$ satisfies descent, it is fibrant in a particular local model structure on presheaves (see [21]). Even though X can be viewed as a stack, it is not cofibrant, and so we need to cofibrantly replace it. Indeed, if X is a (paracompact) manifold with good open cover $\{U_i\}_{i \in I}$, we can replace X by its Čech nerve

$$C(\{U_i\}) := \text{hocolim} \left\{ \coprod_i U_i \rightrightarrows \coprod_{i,j} U_{ij} \rightrightarrows \coprod_{i,j,k} U_{ijk} \rightrightarrows \dots \right\} \quad (2.10)$$

which is both cofibrant and weak equivalent to X [17]. For purely model category theoretic reasons it then follows that $\text{Map}(C(\{U_i\}), \mathbb{B}^n U(1)_{\text{conn}})$ is a Kan complex and we can take π_0 , obtaining the set of morphisms in the homotopy category. This motivates the definition

$$\text{Map}(X, \mathbb{B}^n U(1)_{\text{conn}}) := \text{Map}(C(\{U_i\}), \mathbb{B}^n U(1)_{\text{conn}}) .$$

As explained in [21], these stacks also have a nice geometric interpretation. The following example illustrates the point quite well.

Example 1. Let X be a manifold. Let us calculate the set of vertices of the mapping space $\text{Map}(X, \mathbb{B}^2 U(1)_{\text{conn}})$. Using the pointwise formula for the homotopy colimit [21], we have

$$\begin{aligned} \text{hom}(X, \mathbb{B}^2 U(1)_{\text{conn}}) &= \text{hom}(C(\{U_i\}), \mathbb{B}^2 U(1)_{\text{conn}}) \\ &= \text{hom} \left(\int^{k \in \Delta} \Delta[k] \odot \prod_{\alpha_1, \dots, \alpha_k} U_{\alpha_1, \dots, \alpha_k}, \mathbb{B}^2 U(1)_{\text{conn}} \right) \\ &= \int_{k \in \Delta} \prod_{\alpha_1, \dots, \alpha_k} \text{hom}(\Delta[k] \odot U_{\alpha_1, \dots, \alpha_k}, \mathbb{B}^2 U(1)_{\text{conn}}) \\ &= \prod_{\alpha_1, \dots, \alpha_k} \int_{k \in \Delta} \text{hom}(\Delta[k], \mathbb{B}^2 U(1)_{\text{conn}}(U_{\alpha_1, \dots, \alpha_k})) . \end{aligned} \quad (2.11)$$


An element of the hom in the last line can be written out explicitly as a choice maps

$$\begin{aligned} B_\alpha &: \Delta[0] \rightarrow \prod_{\alpha} \mathbb{B}^2 U(1)(U_\alpha) \\ A_{\alpha\beta} &: \Delta[1] \rightarrow \prod_{\alpha\beta} \mathbb{B}^2 U(1)(U_{\alpha\beta}) \\ g_{\alpha\beta\gamma} &: \Delta[2] \rightarrow \prod_{\alpha\beta\gamma} \mathbb{B}^2 U(1)(U_{\alpha\beta\gamma}) , \end{aligned} \quad (2.12)$$

such that the face inclusions of each map are equal to their corresponding restrictions to higher intersections. Now since equivalent stacks will produce the same cohomology groups, we do not distinguish between equivalent stacks. In particular, using the exponential quasi-isomorphism, we could have equivalently defined $\mathbb{B}^2 U(1)_{\text{conn}}$ to be the stackification of the prestack given by applying the Dold-Kan functor to the presheaf of chain complexes

$$[0 \rightarrow \dots \rightarrow C^\infty(-, U(1)) \xrightarrow{d \log} \Omega^1 \rightarrow \Omega^2] .$$

We can therefore describe the choices of $B_\alpha, A_{\alpha\beta}$ and $g_{\alpha,\beta\gamma}$ via the 2-simplex



Here, $g_{\alpha\beta\gamma}$ is a choice of smooth $U(1)$ -valued function on triple intersections, $A_{\alpha\beta}$ is a choice of 1-form on double intersections and B_α is a choice of 2-form on open sets. Moreover, we have that these assignments must satisfy the conditions

1. $g_{\alpha\beta} g_{\gamma\beta}^{-1} g_{\gamma\alpha} = 1;$
2. $g_{\alpha\beta\gamma}^{-1} dg_{\alpha\beta\gamma} = d\log(g)_{\alpha\beta\gamma} = A_{\alpha\beta} - A_{\gamma\beta} + A_{\gamma\alpha};$
3. $B_\beta - B_\alpha = dA_{\alpha\beta}.$

We identify this data as precisely giving a gerbe with connection [5]. Moreover, the fact that $\mathbb{B}^n U(1)_{\text{conn}}$ is a stack ensures that $F_\alpha = dB_\alpha$ is a globally defined 3-form: the curvature of the gerbe. Notice that these are only the vertices in the mapping space. The entire mapping space keeps track of more information, namely the homotopies and higher homotopies between gerbes. These encode automorphisms in the sense of gauge transformations (see [19] [20]).

Example 2. Let X be a paracompact manifold and $C(\{U_i\})$ the Čech nerve of some good open cover. The maps

$$L : C(\{U_i\}) \rightarrow \mathbb{B}U(1)_{\text{conn}}$$

are in bijective correspondence with circle bundles equipped with a connection. In fact, using the calculations in the above example shows that such a morphism gives the data $U(1)$ -valued functions $g_{\alpha\beta}$ on intersections satisfying $g_{\alpha\beta} g_{\beta\gamma}^{-1} g_{\gamma\delta} = 1$ on triple intersections, along with 1-forms A_α on open sets satisfying $A_\alpha - A_\beta = d\log(g)_{\alpha\beta}$ on double intersections. If the homotopy class of L is trivial, then the circle bundle is trivializable. In fact, the trivializing map ϕ is nothing but a homotopy $\phi : L \rightarrow 0$. To identify this homotopy, we use the Dold-Kan correspondence. In particular, an edge in $\text{Map}(C(\{U_i\}), \mathbb{B}U(1)_{\text{conn}})$ is, by adjunction, an edge in the simplicial set

$$\text{Map}(C(\{U_i\}), \mathbb{B}^n U(1)_{\text{conn}}) = DK(\text{hom}_{\text{Ch}^+}(N(C(\{U_i\})), \mathbb{Z}_{\mathcal{D}}^\infty[2])) , \quad (2.13)$$

where N is the normalized Moore functor. Recall that this functor gives an equivalence of categories, from simplicial abelian groups $s\text{Ab}$ to chain complexes in non-negative degree Ch_\bullet^+ (see [24]). The hom in positively graded chain complexes is the truncated total complex of the Čech-Deligne double complex

$$[\dots \rightarrow \text{tot}^1 C(\mathcal{U}, \mathbb{Z}_{\mathcal{D}}^\infty[2]) \rightarrow Z(\text{tot}^2 C(\mathcal{U}, \mathbb{Z}_{\mathcal{D}}^\infty[2]))] ,$$

where Z denotes the group of cocycles in that degree. Recalling that the differential is given by $D := d + (-1)^k \delta$, where δ takes the alternating sum of restrictions, we identify an edge connecting L and 0 as an assignment of Čech-Deligne cochain h of degree 1 such that $(d - \delta)h = L$. Explicitly, this means a choice of $U(1)$ -valued function h_α on open sets such that

1. $h_\alpha h_\beta^{-1} = g_{\alpha\beta}$;
2. $-ih_\alpha^{-1} dh_\alpha = d \log(h_\alpha) = A_\alpha$.

A straightforward calculation shows that the pattern continues and that null homotopies of n -gerbes (equivalently n -bundles, equivalently maps into $\mathbb{B}^n U(1)_{\text{conn}}$) can again be identified with trivializations.

Motivated by this last example, we will often refer to null homotopies as *trivializations*. To summarize, the mapping space $\text{Map}(X, \mathbb{B}^n U(1)_{\text{conn}})$ can be identified with the set of all n -gerbes with connection, along with isomorphisms between these and higher homotopies between these isomorphisms.

Remark 3. *There are several other stacks related to $\mathbb{B}^n U(1)_{\text{conn}}$ which are useful for us and are defined as follows (see [21],[19],[20],[47]):*

1. *If we forget about the connection on these n -bundles, we obtain the bare moduli stack of n -gerbes $\mathbb{B}^n U(1)$. Explicitly, this stack is obtained by applying the Dold-Kan functor to the sheaf of chain complexes $C^\infty(-, U(1))[n]$: the sheaf of smooth $U(1)$ -valued functions in degree n .*
2. *We also define a stack which represents flat n -bundles with connection, $\flat \mathbb{B}^n U(1)$. This stack is obtained by applying Dold-Kan to the sheaf of chain complexes $\text{disc} U(1)[n]$: the sheaf of locally constant $U(1)$ valued functions in degree n , where “disc” refers to the underlying discrete topology.¹*
3. *We have a stack representing the truncated de Rham complex $\flat_{\text{dR}} \mathbb{B}^n U(1)$ obtained by applying Dold-Kan to the truncated de Rham sheaf of chain complexes*

$$\Omega_{\text{cl}}^{\leq n} := [\dots 0 \dots \Omega^0 \rightarrow \Omega^1 \rightarrow \dots \Omega_{\text{cl}}^n] .$$

4. *Finally, we define the stack of closed n -forms Ω_{cl}^n to be the stack obtained by applying Dold-Kan to the sheaf of closed n -forms.*

One way to see that the second stack really does detect *flat* n -gerbes with connection is to observe that, by Poincaré lemma, one has a quasi-isomorphism of sheaves

$$\text{disc}(U(1))[n] \simeq [0 \rightarrow \dots \rightarrow C^\infty(-, U(1)) \xrightarrow{d \log} \Omega^1 \rightarrow \dots \rightarrow \Omega_{\text{cl}}^n] ,$$

where on the right we have *closed* n -forms in degree 0. These n -forms are to be interpreted as giving the connection on the corresponding bundle. Hence, if the form is closed then the bundle is flat.

The moduli stack $\mathbb{B}^n U(1)_{\text{conn}}$ is related to the stacks in Remark 3 in various ways. In [21][47], it was observed that $\mathbb{B}^n U(1)_{\text{conn}}$ is the homotopy pullback

$$\begin{array}{ccc} \mathbb{B}^n U(1)_{\text{conn}} & \xrightarrow{R} & \Omega_{\text{cl}}^{n+1} \\ \downarrow & & \downarrow \iota \\ \mathbb{B}^n U(1) & \xrightarrow{\theta} & \flat_{\text{dR}} \mathbb{B}^{n+1} U(1) , \end{array} \tag{2.14}$$

¹sometimes alternatively denoted with a superscript “ $(-)^{\delta}$ ”.

where the left composite $\mathbb{B}^n U(1)_{\text{conn}} \rightarrow \mathbb{B}^n U(1) \xrightarrow{\theta} \mathfrak{b}_{\text{dR}} \mathbb{B}^{n+1} U(1)$ is homotopic to the map

$$\text{curv} : \mathbb{B}^n U(1)_{\text{conn}} \rightarrow \mathfrak{b}_{\text{dR}} \mathbb{B}^{n+1} U(1) \quad (2.15)$$

induced by the morphism of sheaves of chain complexes

$$\begin{array}{ccc} \mathbb{Z} & \xrightarrow{i} & \Omega^0 \\ \downarrow i & & \downarrow d \\ \Omega^0 & \xrightarrow{d} & \Omega^1 \\ \downarrow & & \downarrow \\ \vdots & \xrightarrow{d} & \vdots \\ \downarrow & & \downarrow \\ \Omega^n & \xrightarrow{d} & \Omega_{\text{cl}}^{n+1} . \end{array} \quad (2.16)$$

This map gives the full de Rham data for the curvature of a bundle with connection. In fact, since each degree on the right side is a fine sheaf, the sheaf hypercohomology in degree 0 is $H_{\text{dR}}^n(X)$ and the map curv induces a map

$$\text{curv}_* : \pi_0 \text{Map}(X, \mathbb{B}^n U(1)_{\text{conn}}) \rightarrow H_{\text{dR}}^{n+1}(X) , \quad (2.17)$$

which sends an n -bundle with connection to the de Rham class of its curvature. The following proposition might certainly be known to experts, but we include a proof for completeness.

Lemma 2. *The homotopy fiber of the map*

$$R : \mathbb{B}^n U(1)_{\text{conn}} \rightrightarrows \Omega_{\text{cl}}^{n+1}$$

can be identified with $\mathfrak{b} \mathbb{B}^n U(1)$.

Proof. The map R is induced by the morphism of sheaves of chain complexes

$$\begin{array}{ccc} \underline{\mathbb{Z}} & \xrightarrow{0} & 0 \\ \downarrow i & & \downarrow d \\ \Omega^0 & \xrightarrow{0} & 0 \\ \downarrow & & \downarrow \\ \vdots & \xrightarrow{0} & \vdots \\ \downarrow & & \downarrow \\ \Omega^{n-1} & \xrightarrow{d} & \Omega_{\text{cl}}^n . \end{array} \quad (2.18)$$

Since this map is degree-wise surjective by Poincaré lemma (traditionally in highest form-degree, and trivially in lower degrees), it is a fibration in the projective model structure on presheaves of chain complexes. We can therefore calculate the homotopy fiber as the kernel of that map. By inspection, the kernel is

$$[\dots \rightarrow \mathbb{Z} \hookrightarrow \Omega^0 \rightarrow \Omega^1 \rightarrow \dots \Omega_{\text{cl}}^n],$$

which, via the exponential map, is quasi-isomorphic to

$$[\dots C^\infty(-, U(1)) \xrightarrow{d\log} \Omega^1 \rightarrow \dots \Omega_{\text{cl}}^n].$$

Again, by Poincaré lemma, this sheaf of chain complex is quasi-isomorphic to $\text{disc}(U(1))[n]$. Since the Dold-Kan functor is a right Quillen adjoint and preserves weak equivalences, it takes fibration sequences to fibration sequences and we have the desired result. \square

Using the above proposition along with diagram (2.14) and the pasting lemma for homotopy pullbacks, we observe that we have the following iteration of homotopy pullbacks [47]

$$\begin{array}{ccccccc}
\mathbb{b}\mathbb{B}^{n-1}U(1) & \longrightarrow & \mathbb{B}^nU(1) & \longrightarrow & * & & \\
\downarrow & & \downarrow & & \downarrow & & \\
* & \longrightarrow & \mathbb{b}_{\text{dR}}\mathbb{B}^{n-1}U(1) & \longrightarrow & \mathbb{b}\mathbb{B}^nU(1) & \longrightarrow & * \\
\downarrow & & \downarrow & & \downarrow & & \downarrow \\
* & \longrightarrow & \Omega^{\leq n-1} & \longrightarrow & \mathbb{B}^nU(1)_{\text{conn}} & \twoheadrightarrow & \Omega_{\text{cl}}^{n+1} \\
& & \downarrow & & \downarrow & & \downarrow \\
& & * & \longrightarrow & \mathbb{B}^nU(1) & \longrightarrow & \mathbb{b}_{\text{dR}}\mathbb{B}^{n+1}U(1).
\end{array} \tag{2.19}$$

From Lemma 2 along with this last diagram, we immediately get the following:

Proposition 3. *The loop stack $\Omega\mathbb{B}^nU(1)_{\text{conn}}$ can be identified with the stack $\mathbb{b}\mathbb{B}^{n-1}U(1)$.*

Proof. Consider the homotopy pullback square

$$\begin{array}{ccccc}
\mathbb{b}\mathbb{B}^{n-1}U(1) & \longrightarrow & \mathbb{B}^nU(1) & \longrightarrow & * \\
\downarrow & & \downarrow & & \downarrow \\
* & \longrightarrow & \mathbb{b}_{\text{dR}}\mathbb{B}^{n-1}U(1) & \longrightarrow & \mathbb{b}\mathbb{B}^nU(1) \\
\downarrow & & \downarrow & & \downarrow \\
* & \longrightarrow & \Omega^{\leq n-1} & \longrightarrow & \mathbb{B}^nU(1)_{\text{conn}}
\end{array}$$

within diagram (2.19). Such a homotopy pullback, given by the outer square, can be taken as a definition of the loop space. Alternatively, a homotopy pullback can be computed explicitly as the paths in $\mathbb{B}^n U(1)_{\text{conn}}$ connecting the point inclusion $* \rightarrow \mathbb{B}^n U(1)_{\text{conn}}$ to itself: a loop. \square

Note that Massey products in the homology of the free loop space is classically considered in [46] [11]. The above discussions allows us to recast the ‘differential cohomology diamond’ using our stacks.

Proposition 4. *The differential cohomology diagram (2.7) lifts to a diagram of stacks*

$$\begin{array}{ccccc}
 & \Omega^{\leq n-1} & \xrightarrow{d} & \Omega_{\text{cl}}^n & \\
 & \swarrow a & & \nearrow R & \\
 {}_{\text{b}dR}\mathbb{B}^{n-1}U(1) & & \mathbb{B}^n U(1)_{\text{conn}} & & {}_{\text{b}dR}\mathbb{B}^n U(1) \\
 & \searrow & \nearrow I & & \swarrow \\
 {}_{\text{b}}\mathbb{B}^n U(1) & \xrightarrow{\quad} & \mathbb{B}^n U(1) & &
 \end{array} \tag{2.20}$$

where the diagonals are fibration sequences.

Proof. This is the same diagram as a portion of diagram (2.19) rotated. \square

We now explain how to go the other direction, i.e. from stacks to Deligne cohomology. We have seen that for a manifold X , the mapping space $\text{Map}(X, \mathbb{B}^n U(1)_{\text{conn}})$ can be identified with the space of n -gerbes equipped with connections (along with all isomorphisms and higher isomorphisms between them). It will be convenient to organize this mapping space itself into a stack. We define the *mapping stack* to be the stackification of the prestack given by the assignment

$$U \mapsto \text{Map}(X \times U, \mathbb{B}^n U(1)_{\text{conn}}) \tag{2.21}$$

for each $U \in \text{CartSp}$. We denote this stack by $^2[X, \mathbb{B}^n U(1)_{\text{conn}}]$.

Remark 4. *Notice the following:*

1. *If we evaluate the mapping stack on the terminal object in CartSp (the point) and take π_0 , we recover the usual differential cohomology groups*

$$\pi_0[X, \mathbb{B}^n U(1)_{\text{conn}}](*) \simeq \pi_0 \text{Map}(X \times *, \mathbb{B}^n U(1)_{\text{conn}}) \simeq \widehat{H}^n(X, \mathbb{Z}) .$$

2. *Since the mapping stack is clearly functorial in both arguments and the stackification functor preserves homotopy fibers (it is left exact), we can map into the diagram (2.19) to obtain the*

²Note that this is not to be confused with homotopy classes of maps as the notation might suggest.

diagram

$$\begin{array}{ccccc}
& [X, \Omega^{\leq n-1}] & \xrightarrow{d} & [X, \Omega_{cl}^n] & \\
& \nearrow & \searrow a & \nearrow R & \searrow \\
[X, \flat_{dR} \mathbb{B}^{n-1} U(1)] & & [X, \mathbb{B}^n U(1)_{\text{conn}}] & & [X, \flat_{dR} \mathbb{B}^n U(1)] \\
& \searrow & \nearrow & \searrow I & \nearrow \\
& [X, \flat \mathbb{B}^n U(1)] & \xrightarrow{\quad} & [X, \mathbb{B}^n U(1)] &
\end{array}$$

where the diagonals are again fibration sequences. If we evaluate this previous diagram at the point and apply π_0 , we indeed reproduce the usual differential cohomology diagram (2.19).

2.3 Cup product in differential cohomology

Deligne [13] and Beilinson [3] showed that differential cohomology admits a distinguished cup product refining the usual cup product on singular cohomology. This product is defined on sections of $\mathbb{Z}_{\mathcal{D}}^{\infty}[n]$ by the formula

$$\alpha \cup_{DB} \beta = \begin{cases} \alpha \beta & \deg(\alpha) = n \\ \alpha \wedge d\beta & \deg(\alpha) = 0 \\ 0 & \text{otherwise.} \end{cases} \quad (2.22)$$

Note that the grading here is such that the first case is simply multiplication by an integer. In fact, it is obvious from the definition that the Deligne-Beilinson (henceforth DB) cup product composed with the natural inclusion

$$\mathbb{Z}[n] \hookrightarrow \mathbb{Z}_{\mathcal{D}}^{\infty}[n]$$

simply multiplies the two locally constant integer-valued functions. Since the sheaf cohomology of the locally constant sheaf \mathbb{Z} , equipped with this product, is simply the ordinary cohomology ring with integral coefficients, one immediately sees that this cup product does indeed refine the usual cup product.

Equipped with this cup product, $\widehat{H}^*(X; \mathbb{Z})$ becomes an associative and graded-commutative ring [5]. This cup product structure also refines the wedge product of forms in the sense that the curvature map $R : \widehat{H}^*(X; \mathbb{Z}) \rightarrow \Omega_{cl}^*$ defines a homomorphism of graded commutative rings [6]. In particular this implies that the cup product of two classes of odd degree is flat. It can also be shown [6] that the cup product of a flat class with any other class is again flat and that the inclusion of $H^*(X, U(1))$ into $\widehat{H}^*(X; \mathbb{Z})$ is a two sided ideal.

We now turn to the cup product, viewed as a morphism of stacks. In [19] it was observed that the lax monoidal structure of the Dold-Kan map gives rise to a cup product, exhibited as a morphism

$$\cup : \mathbb{B}^m U(1)_{\text{conn}} \times \mathbb{B}^n U(1)_{\text{conn}} \rightarrow \mathbb{B}^{n+m+1} U(1)_{\text{conn}} \quad (2.23)$$

of stacks. This map is obtained by simply taking the DB cup product (2.22)

$$\cup_{DB} : \mathbb{Z}_{\mathcal{D}}^{\infty}[n+1] \otimes \mathbb{Z}_{\mathcal{D}}^{\infty}[m+1] \rightarrow \mathbb{Z}_{\mathcal{D}}^{\infty}[n+m+2],$$

applying the Dold-Kan map

$$DK(\cup_{DB}) : DK(\mathbb{Z}_{\mathcal{D}}^{\infty}[n+1] \otimes \mathbb{Z}_{\mathcal{D}}^{\infty}[m+1]) \rightarrow DK(\mathbb{Z}_{\mathcal{D}}^{\infty}[n+m+2]) ,$$

and using the lax monoidal structure φ of DK to get a map

$$\cup = DK(\cup_{DB}) \circ \varphi : DK(\mathbb{Z}_{\mathcal{D}}^{\infty}[n+1]) \times DK(\mathbb{Z}_{\mathcal{D}}^{\infty}[m+1]) \rightarrow DK(\mathbb{Z}_{\mathcal{D}}^{\infty}[n+1] \otimes \mathbb{Z}_{\mathcal{D}}^{\infty}[m+1]) \rightarrow DK(\mathbb{Z}_{\mathcal{D}}^{\infty}[n+m+2]) .$$

Applying the stackification functor then gives the desired map. This map then induces a map of stacks (which we also denote as \cup)

$$\cup : [X, \mathbb{B}^n U(1)_{\text{conn}}] \times [X, \mathbb{B}^m U(1)_{\text{conn}}] \rightarrow [X, \mathbb{B}^{n+m+1} U(1)_{\text{conn}}] . \quad (2.24)$$

The following two propositions are implicit in [19] [20].

Proposition 5. *The DB cup product refines the singular cup product. That is, we have a commutative diagram*

$$\begin{array}{ccc} \mathbb{B}^n U(1)_{\text{conn}} \times \mathbb{B}^m U(1)_{\text{conn}} & \xrightarrow{\cup_{DB}} & \mathbb{B}^{n+m+1} U(1)_{\text{conn}} \\ \downarrow I \times I & & \downarrow I \\ \mathbb{B}^{n+1} \mathbb{Z} \times \mathbb{B}^{m+1} \mathbb{Z} & \xrightarrow{\cup} & \mathbb{B}^{n+m+2} \mathbb{Z} . \end{array}$$

Proof. Let $p : \mathbb{Z}_{\mathcal{D}}^{\infty}[n+1] \rightarrow \mathbb{Z}[n+1]$ be the projection map

$$\begin{array}{ccc} \mathbb{Z} & \xrightarrow{\text{id}} & \mathbb{Z} \\ \downarrow i & & \downarrow d \\ \Omega^0 & \xrightarrow{0} & 0 \\ \downarrow & & \downarrow \\ \vdots & \xrightarrow{0} & \vdots \\ \downarrow & & \downarrow \\ \Omega^n & \xrightarrow{0} & 0 . \end{array}$$

Then by definition of the DB cup product, the diagram

$$\begin{array}{ccc} \mathbb{Z}_{\mathcal{D}}^{\infty}[n+1] \otimes \mathbb{Z}_{\mathcal{D}}^{\infty}[m+1] & \xrightarrow{\cup_{DB}} & \mathbb{Z}_{\mathcal{D}}^{\infty}[n+m+2] \\ \downarrow p & & \downarrow p \\ \mathbb{Z}[n+1] \otimes \mathbb{Z}[m+1] & \xrightarrow{\cup} & \mathbb{Z}[n+m+2] \end{array}$$

commutes in sheaves of chain complexes. Applying the Dold-Kan functor and using naturality of the lax monoidal structure map gives the result. \square

Proposition 6. *The cup product refines the wedge product, and we have a commutative diagram*

$$\begin{array}{ccc} \mathbb{B}^n U(1)_{\text{conn}} \times \mathbb{B}^m U(1)_{\text{conn}} & \xrightarrow{\cup_{DB}} & \mathbb{B}^{n+m+1} U(1)_{\text{conn}} \\ \downarrow R \times R & & \downarrow R \\ \Omega_{\text{cl}}^{n+1} \times \Omega_{\text{cl}}^{m+1} & \xrightarrow{\wedge} & \Omega_{\text{cl}}^{n+m+2} . \end{array}$$

Proof. Let α and β be sections of $\mathbb{Z}_{\mathcal{D}}^{\infty}[n+1]$ and $\mathbb{Z}_{\mathcal{D}}^{\infty}[m+1]$, respectively. Applying the curvature R to the DB cup product (2.22) gives

$$R(\alpha \cup_{DB} \beta) = \begin{cases} \alpha d(\beta) & \text{if } \deg(\alpha) = n \\ d(\alpha) \wedge d(\beta) & \text{if } \deg(\beta) = 0 \\ 0 & \text{otherwise} \end{cases}$$

which is $R(\alpha) \wedge R(\beta)$. We therefore have a commuting diagram

$$\begin{array}{ccc} \mathbb{Z}_{\mathcal{D}}^{\infty}[n+1] \otimes \mathbb{Z}_{\mathcal{D}}^{\infty}[m+1] & \xrightarrow{\cup_{DB}} & \mathbb{Z}_{\mathcal{D}}^{\infty}[n+m+2] \\ \downarrow R & & \downarrow R \\ \Omega_{\text{cl}}^{n+1} \otimes \Omega_{\text{cl}}^{m+1} & \xrightarrow{\wedge} & \Omega_{\text{cl}}^{n+m+2} , \end{array}$$

Applying Dold-Kan gives the result in stacks. \square

The above results show that, in general, the Deligne-Beilinson cup product does not refine the de Rham wedge product for the whole de Rham complex, but does so only for the top and bottom degrees. However, for the triple product the only cup products that arise are between degree zero and degree one cocycles, so that nothing is missed in passing to \cup_{DB} . We will make this more precise in Prop. 17.

2.4 Massey products in hypercohomology

Massey products in Deligne-Beilinson cohomology are described in [16] [44] [48] [38]. In this section, we review the construction for hypercohomology found in [44], with a slightly adapted language for later comparison and generalization. In section 3 we generalize this construction in two ways, which we describe. We use the Dold-Kan correspondence to establish these products in the stacky setting. We also use the machinery of May [35] to exhibit these products as differential matrix Massey products.

Let R be a commutative ring and let $\mathcal{C}^{\bullet}(n)$, $n \in \mathbb{N}$, be a sequence of positively graded chain complexes of R -modules. Moreover, let us assume that this sequence comes equipped with maps

$$\cup : \mathcal{C}^{\bullet}(n) \otimes \mathcal{C}^{\bullet}(m) \rightarrow \mathcal{C}^{\bullet}(n+m) ,$$

which are associative in the sense that

$$\cup \circ (id \otimes \cup) = \cup \circ (\cup \otimes id) . \quad (2.25)$$

The maps \cup induce an associative product on cohomology

$$\cup : H^\bullet(n) \otimes H^\bullet(m) \rightarrow H^\bullet(n+m) ,$$

called the *cup product*. Once a well-defined notion of a cup product is established, one can define the Massey products via the following.

Definition 7. Let $l \geq 2$ and let n_1, \dots, n_l and m_1, \dots, m_l be integers. Define

$$n_{s,t} = \sum_{i=s}^t (n_i - 1) \quad \text{and} \quad m_{s,t} = \sum_{i=s}^t m_i, \quad \text{for } 1 \leq s \leq t \leq l ,$$

and let $\bar{a} = (-1)^{q+1}a$ denote the twist of a class $a \in \mathcal{C}^q(n)$. We define the l -fold Massey product as follows:

(i). Let $a_i \in H^{m_i}(\mathcal{C}^\bullet(n_i))$ be cohomology classes. Suppose there exists cochains $a_{s,t} \in \mathcal{C}^{m_{s,t}+1}(n_{s,t})$ such that $a_{i,i}$ is a representative of a_i and that

$$da_{s,t} = \sum_{i=s}^{t-1} \bar{a}_{s,i} \cup a_{i+1,t} \quad \text{for } 1 \leq s \leq t \leq l, \quad (s,t) \neq (1,l) .$$

We call the collection $\mathcal{M} = \{a_{s,t}\}$ a defining system for the l -fold Massey product.

(ii). The cochain

$$a_{1,l} := \sum_{i=1}^{l-1} \bar{a}_{1,i} \cup a_{i+1,l} \in \mathcal{C}^{m_{1,l}+2}(n_{1,l})$$

is a cocycle and represents a cohomology class m_l . We call this class the l -fold Massey product of the elements a_1, \dots, a_l with defining system \mathcal{M} .

In general, we would like to eliminate the dependance of the product on the defining system. The case of $l = 3$ will be the most important for us, and in this case we are indeed able to eliminate this dependence. The following three examples are known, and we record them to highlight how Massey products arise in the different settings that we consider, and how stacks will provide, in a sense, a unifying theme. Note that, while the above construction is fairly general, it is not obvious how to generalize to other settings and how to do computations easily with it, and that is why we later use the stacky perspective.

Example 3. Let a_1, a_2 and a_3 be cohomology classes as above. Suppose we have a defining system $\mathcal{M} = \{a_{s,t}\}$. This means, by definition, that we have the relations

$$da_{1,2} = \bar{a}_{1,1} \cup a_{2,2} \quad \text{and} \quad da_{2,3} = \bar{a}_{2,2} \cup a_{3,3} .$$

Now a class m_l representing the Massey product of this defining system has as a representing cocycle

$$a_{1,l} = \bar{a}_{1,1} \cup a_{2,3} + \bar{a}_{1,2} \cup a_{3,3} .$$

Notice that, in this case, the class m_l only depends on the defining system up to cocycles. That is, for another defining system $\mathcal{N} = \{b_{s,t}\}$, the classes $a_{2,3} - b_{2,3}$ and $a_{1,2} - b_{1,2}$ are cocycles. Moreover, if these cocycles are coboundaries, then the Massey products of both defining systems agree. We can therefore define a Massey product, not depending on the defining system, as the quotient

$$\langle a_1, a_2, a_3 \rangle \in \frac{H^{m_1+m_2+m_3-1}(\mathcal{C}^\bullet(n_1+n_2+n_3))}{H^{m_1+m_2-1}(\mathcal{C}^\bullet(n_1+n_2)) \cup a_3 + a_1 \cup H^{m_2+m_3-1}(\mathcal{C}^\bullet(n_2+n_3))} .$$

Example 4. Let X be a smooth manifold and let $C(n) = \Omega^*(X)$ for each n , where $\Omega^*(X)$ is the algebra of differential forms on X . Let a, b and c be de Rham cohomology classes of degree p, q, r respectively, such that $a \wedge b = 0 = b \wedge c$. Choose representing closed forms α, β, γ for a, b, c respectively, and let η and ρ be cochains such that

$$d\eta = \alpha \wedge \beta \quad \text{and} \quad d\rho = \beta \wedge \gamma .$$

Then the combination

$$\eta \wedge \gamma - (-)^p \alpha \wedge \rho$$

is a closed form representing the triple Massey product of a, b and c corresponding to the defining system $\mathcal{M} = (\alpha, \beta, \gamma, \rho, \eta)$. Eliminating the dependence on \mathcal{M} gives a well-defined class in the quotient group

$$H_{\text{dR}}^{p+q+r-1}(X) / \left(a \cup H_{\text{dR}}^{q+r-1}(X) + c \cup H_{\text{dR}}^{p+q-1}(X) \right) .$$

The following constitutes our initial transition to differential cohomology, which we will develop in stacks in the following section.

Example 5. Consider the Deligne complex given by the sheaf of chain complexes

$$\mathbb{Z}_{\mathcal{D}}^\infty[n] := [\underline{\mathbb{Z}} \hookrightarrow \Omega^0 \rightarrow \Omega^1 \rightarrow \dots \rightarrow \Omega^{n-1}] .$$

Let X be a paracompact manifold with good open cover $\{U_i\}_{i \in I}$ and let $C(n) := \text{tot} C^\bullet(\{U_i\}, \mathbb{Z}_{\mathcal{D}}^\infty[n])$ be the total complex of the Čech-Deligne double complex. The degree n cohomology of this total complex calculates the differential cohomology of X :

$$\hat{H}^n(X; \mathbb{Z}) = H^n \text{tot} C^\bullet(\mathcal{U}, \mathbb{Z}_{\mathcal{D}}^\infty[n]) .$$

The Deligne-Beilinson cup product is defined as a morphism

$$\cup_{DB} : \mathbb{Z}_{\mathcal{D}}^\infty[n] \otimes \mathbb{Z}_{\mathcal{D}}^\infty[m] \rightarrow \mathbb{Z}_{\mathcal{D}}^\infty[n+m+1] ,$$

which on sections is given by the formula (2.22). This map induces cup product morphisms on the total complexes $C(n)$ which are associative in the sense of the identity in (2.25).

We can therefore use this cup product to define the Massey product in differential cohomology, viewed as the sheaf hypercohomology of the Deligne complex. Since our point of view will subsume this construction, we will delay explicit examples until sections 3 and 4.

3 Massey products in the language of higher stacks

We provide our main construction of stacky Massey products in this section. We start with setting up the machinery needed.

3.1 The Dold-Kan correspondence

The Dold-Kan correspondence will be an important component in defining the Massey product in stacks. We will use the correspondence to organize the homotopies involved in certain homotopy commuting diagrams in an algebraic way.

The classical Dold-Kan correspondence describes an equivalence of categories (see e.g. [24])

$$\Gamma : \mathcal{Ch}^+ \xrightleftharpoons{\quad} s\mathcal{Ab} : N \quad (3.1)$$

between positively graded chain complexes and simplicial abelian groups. By post-composing with the free-forgetful adjunction, one obtains an adjunction

$$DK := UT : \mathcal{Ch}^+ \xrightleftharpoons{\quad} s\mathcal{Set} : NF. \quad (3.2)$$

In fact, one can say more. This adjunction is a Quillen adjunction of model categories, with the projective model structure on chain complexes and the Quillen model structure on simplicial sets. As such, it preserves the homotopy theories in both categories; it therefore comes as no surprise that for a positively graded chain complex C_\bullet one has an isomorphism

$$H_n C_\bullet \simeq \pi_n DK(C_\bullet). \quad (3.3)$$

For convenience, we remind the reader what the functor DK does to a chain complex, as this will be a frequently used tool in producing abelian stacks.

Let Δ denote the category of linearly ordered sets of n elements with order preserving maps. Let C_\bullet be a positively graded chain complex. The degree n component of the simplicial abelian group $DK(C_\bullet)$ is given by

$$DK(C_\bullet)_n = \bigoplus_{[n] \twoheadrightarrow [k]} C_k.$$

Here the indexing set is taken to be all *surjections* $[n] \twoheadrightarrow [k]$. It is a bit trickier to describe the face and degeneracy maps. Let $d^i : [n-1] \hookrightarrow [n]$ be a coface map in Δ . We want to define the corresponding face map. To get a map out of the direct sum, it suffices to describe the map on each factor. Therefore, we need only define the face map on a term C_k given by a surjection $\sigma : [n] \twoheadrightarrow [k]$. To see where to send this term, we form the composite $\sigma d^i[n-1] \hookrightarrow [n] \twoheadrightarrow [k]$. Now this morphism need not be surjective, so we factorize $\mu \sigma'[n-1] \twoheadrightarrow [m] \hookrightarrow [k]$ where the first map is a surjection and the second map is an injection. Then σ' corresponds to a term $C_m \hookrightarrow \bigoplus_{[n-1] \twoheadrightarrow [m]} A_m = DK(C_\bullet)_{n-1}$. We send the factor C_k to the factor C_m by a map $\mu' : C_k \rightarrow C_m$. This map is given by

$$\mu' = \begin{cases} id & \mu = id \\ (-1)^k d & \mu = d^k \\ 0 & \text{otherwise.} \end{cases} \quad (3.4)$$

A similar construction is used to define the degeneracy maps. The following example illustrates the point quite well.

Example 6. Consider the chain complex $A[1]$, with the abelian group A in degree 1 and 0's in all other degrees. We want to compute $DK(A[1])$. Using the above formula, we see that the only nonzero terms in degree n are given by the surjections $[n] \twoheadrightarrow [1]$. Each surjection can be thought of as being given by an element $i \in [n]$ which divides the set into two subsets: those that go to 0 and those that go to 1. We therefore have n surjections and


$$DK(A[1])_n = \bigoplus_{i=1}^n A.$$

For a coface map $d^j : [n-1] \rightarrow [n]$, the corresponding face map d^j is given as follows. Let A_i denote the copy of A corresponding to the i th surjection. Then

$$d_j(A_i) = \begin{cases} A_{i-1} & \text{if } i > j \neq 0, n \\ A_i & \text{if } i \leq j \neq 0, n \end{cases}, \quad d_0(A_i) = \begin{cases} A_{i-1} & \text{if } i \neq 0 \\ 0 & \text{if } i = 0 \end{cases}, \quad d_n(A_i) = \begin{cases} A_i & \text{if } i \neq n \\ 0 & \text{if } i = n. \end{cases}$$

Notice that for $j \neq 0, n$, the term corresponding to $i = j$ and $i = j + 1$ both go to the same copy of A . We therefore have a map $A \times A \rightarrow A$ extending the identity on each component. Hence, this morphism is just group multiplication. From this, one can see that this simplicial abelian group is just the delooping group BA .

Another way to describe the simplicial set $DK(C_\bullet)$, which is perhaps more conceptual, is via a labeling of simplices with elements of the chain complex C^\bullet . A 2-simplex in $DK(C^\bullet)$, for example, is a simplex with face, edges and vertices labeled by elements of C^\bullet



such that

1. $dc_{012} = b_{01} + b_{12} - b_{02}$,
2. $db_{ij} = a_j - a_i$.

Here d is the chain complex differential. Notice that a 2-simplex in $DK(C_\bullet)$, defined as before, can be identified as such a labeled simplex. To see this, let us calculate the data involved in specifying a 2-simplex. First, observe that there is exactly one surjection $0 : [2] \twoheadrightarrow [0]$, $\text{id} : [2] \twoheadrightarrow [2]$, and exactly two surjections $\sigma_i : [2] \twoheadrightarrow [1]$. Therefore, a 2-simplex is given by a quadruple $(a_0, b_{01}, b_{02}, c_{012})$, where a_0 is in degree 1, while b_{01}, b_{02} are in degree 2 (corresponding to σ_1, σ_2 resp.), and c_{012} is in degree 3. To determine the edges, we evaluate d_i on this quadruple. For $i = 0$, we have the following epi-mono factorizations

$$\text{id} \circ d^0 = d^0 \circ \text{id}, \quad \sigma_1 \circ d^0 = d^0 \circ 0, \quad \sigma_2 \circ d^0 = \text{id} \circ \text{id}. \quad (3.5)$$

It follows from the formula, that the 0 face is b_{02} . For $i = 1$, we have

$$\text{id} \circ d^1 = d^1 \circ \text{id}, \quad \sigma_1 \circ d^1 = \text{id} \circ \text{id}, \quad \sigma_2 \circ d^1 = \text{id} \circ \text{id} \quad (3.6)$$

and the 1 face is $b_{01} + b_{02}$. Finally, for $i = 2$, we have

$$\text{id} \circ d^2 = d^2 \circ \text{id}, \quad \sigma_1 \circ d^2 = \text{id} \circ \text{id}, \quad \sigma_2 \circ d^2 = d^1 \circ 0 \quad (3.7)$$

and the 2 edge is $dc_{012} + b_{01}$. Forming the boundary of the simplex, we get

$$\partial(a_0, b_{01}, b_{02}, c_{012}) = b_{02} - (b_{01} + b_{02}) + (dc_{012} + b_{01}) = dc_{012}.$$

That the edges of the simplex satisfy the second condition above is a straightforward calculation and will be omitted. In fact, it is a straightforward calculation to show that the boundary of a general n -simplex must be equal to d applied to the labeling on its n -face.

Remark 5. *This second description provides a powerful conceptual advantage; namely, that the differential of the chain complex can be viewed as obstructing the chain from being a cycle. For example, if the resulting simplicial set were the nerve of a groupoid, then all simplices for $n \geq 2$ would be cycles.*

3.2 Stacky Massey products

We are now ready to define Massey products in the category of stacks. We begin with a discussion on Massey triple products and then generalize to l -fold Massey products.

The Massey triple product can be viewed as a homotopy built out of the associativity diagram of the cup product of three elements. In fact, suppose one is given a triple of higher gerbes with connection on a manifold X . These gerbes are given by the data $\mathcal{G}_i : X \rightarrow \mathbb{B}^{n_i}U(1)$, $i = 1, 2, 3$. Suppose, moreover, that these gerbes are chosen so that the cup products $\mathcal{G}_1 \cup \mathcal{G}_2$ and $\mathcal{G}_2 \cup \mathcal{G}_3$ are homotopic to 0, with trivializing homotopies $\phi_{1,2}$ and $\phi_{2,3}$. In this case, we can build a *loop* trivializing the triple product. To see this, consider the associativity diagram for the cup product

$$\begin{array}{ccccc}
& & & \mathbb{B}^{n_1+n_2+1}U(1)_{\text{conn}} \times \mathbb{B}^{n_3}U(1)_{\text{conn}} & \\
& \nearrow & & \downarrow (\cup \times \text{id}) & \searrow \cup \\
X & \xrightarrow{\mathcal{G}_1 \times \mathcal{G}_2 \times \mathcal{G}_3} & \mathbb{B}^{n_1}U(1)_{\text{conn}} \times \mathbb{B}^{n_2}U(1)_{\text{conn}} \times \mathbb{B}^{n_3}U(1)_{\text{conn}} & & \mathbb{B}^{n_1+n_2+n_3+2}U(1)_{\text{conn}} \\
& \searrow & \downarrow (\text{id} \times \cup) & \nearrow \cup & \\
& & \mathbb{B}^{n_1}U(1)_{\text{conn}} \times \mathbb{B}^{n_2+n_3+1}U(1)_{\text{conn}} & &
\end{array}$$

Although the outer two maps agree, there is still nontrivial homotopy theoretic information contained in the diagram. To see this, suppose $\mathcal{G}_1 \cup \mathcal{G}_2$ and $\mathcal{G}_2 \cup \mathcal{G}_3$ are trivializable with trivializations $\phi_{1,2}$ and $\phi_{2,3}$, respectively. Then we can add these homotopies to the diagram

These two choices of homotopies $\phi_{1,2}$ and $\phi_{2,3}$ make the entire diagram homotopy commutative, as the triple cup products (the two red arrows) are trivialized by the homotopies $\mathcal{G}_1 \cup \phi_{1,2}$ and $\phi_{2,3} \cup \mathcal{G}_3$. Since the cup product is strictly associative, the diagram in red commutes and we have the homotopy commuting diagram

$$\begin{array}{ccc}
 & 0 & \\
 & \Downarrow \mathcal{G}_1 \cup \phi_{2,3} & \\
 X & \xrightarrow{\mathcal{G}_1 \cup \mathcal{G}_2 \cup \mathcal{G}_3} & \mathbb{B}^{n_1+n_2+n_3+2}U(1)_{\text{conn}} \\
 & \Uparrow \phi_{1,2} \cup \mathcal{G}_3 & \\
 & 0 &
 \end{array} \quad (3.8)$$

These two homotopies fit together to form a loop. Then, by Prop. 3 and the universal property of the homotopy pullback, we can equivalently describe this as a map

$$X \rightarrow \Omega \mathbb{B}^{n_1+n_2+n_3+2}U(1)_{\text{conn}} \simeq \mathbb{B}^{n_1+n_2+n_3+1}U(1). \quad (3.9)$$

Lemma 8. *The homotopy class of the loop (3.9) is in the image of the inclusion of $H^{n_1+n_2+n_3+1}(X; U(1))$ into $\hat{H}^{n_1+n_2+n_3+2}(X; \mathbb{Z})$.*

Proof. Using the Dold-Kan adjunction along with a Čech resolution of X , we have the following

sequence of isomorphisms

$$\begin{aligned}
\pi_0 \text{Map}(X, {}^b\mathbb{B}^{n_1+n_2+n_3+1}U(1)) &\simeq H_0 \text{hom}_{\mathcal{C}h+}(N(C(\{U_i\}), \text{disc}(U(1)))[n_1 + n_2 + n_3 + 1]) \\
&\simeq H^{n_1+n_2+n_3+1}(X; U(1)) \\
&\hookrightarrow \hat{H}^{n_1+n_2+n_3+2}(X; \mathbb{Z}) .
\end{aligned}$$

Here disc indicates that we are taking the discrete topology on $U(1)$ (i.e. this is the sheaf of locally constant $U(1)$ valued functions). \square

Remark 6. (i) Notice that we could have equivalently taken the homotopy class of the loop directly to get an element $[\mathcal{G}_1 \cup \phi_{2,3} - \phi_{1,2} \cup \mathcal{G}_1]$ in

$$\begin{aligned}
\pi_1 \text{Map}(X, \mathbb{B}^{n_1+n_2+n_3+2}U(1)_{\text{conn}}) &\simeq H_1 \text{hom}_{\mathcal{C}h+}(C(\{U_i\}), \mathbb{Z}_{\mathcal{D}}^{\infty}[n_1 + n_2 + n_3 + 2]) \\
&\simeq H^{n_1+n_2+n_3+1}(X; U(1)) \\
&\hookrightarrow \hat{H}^{n_1+n_2+n_3+2}(X; \mathbb{Z}) .
\end{aligned}$$

(ii) The above observations allow us to recover the usual definition of the Massey product as an element in cohomology. In section 2.4, we observed that such a class is not completely well-defined purely at the level of cohomology and there was some dependence on the chosen cochain representatives. Taking this point of view, one can see this dependence as a choice of trivializations $\phi_{1,2}$ and $\phi_{2,3}$ of the cup products.

This definition works well for the triple product and gives a clear picture on how the triple product is built out of the homotopies. However, to describe the higher triple products this way would be cumbersome. Moreover, the algebraic nature of the products would not be transparent. For these reasons, we will use the language of simplicial homotopy theory to describe these homotopy commuting diagrams and the Dold-Kan correspondence to organize these homotopies in an algebraic way. To prepare the reader for this perspective, we first recast the triple product in this language.

Notice that the triple product was described by two homotopies $\phi_{1,2}$ and $\phi_{2,3}$ connecting the basepoint 0 to the double cup products. We can express this situation diagrammatically via the horn-fillers

$$\begin{array}{ccc}
\partial\Delta[1] & \xrightarrow{(0, \mathcal{G}_1 \cup \mathcal{G}_2)} & [X, \mathbb{B}^{n_1+n_2+1}U(1)_{\text{conn}}] \\
\downarrow & \nearrow \phi_{1,2} & \\
\Delta[1] & &
\end{array}
, \quad
\begin{array}{ccc}
\partial\Delta[1] & \xrightarrow{(0, \mathcal{G}_2 \cup \mathcal{G}_3)} & [X, \mathbb{B}^{n_2+n_3+1}U(1)_{\text{conn}}] \\
\downarrow & \nearrow \phi_{2,3} & \\
\Delta[1] & &
\end{array} .$$

Now we would like to use these homotopies to construct a loop. To do this, we need to manipulate algebraically these homotopies. This motivates us to take the Moore complex of these diagrams in order to translate the data into the language of sheaves of chain complexes. This gives the data

$$\begin{array}{ccc}
\mathbb{Z} \oplus \mathbb{Z} & \xrightarrow{(0, \mathcal{G}_1 \cup \mathcal{G}_2)} & N([X, \mathbb{B}^{n_1+n_2+1}U(1)_{\text{conn}}])_0 \\
\uparrow (1, -1) & & \uparrow \partial \\
\mathbb{Z} & \xrightarrow{\phi_{1,2}} & N([X, \mathbb{B}^{n_1+n_2+1}U(1)_{\text{conn}}])_1
\end{array}
, \quad
\begin{array}{ccc}
\mathbb{Z} \oplus \mathbb{Z} & \xrightarrow{(0, \mathcal{G}_2 \cup \mathcal{G}_3)} & N([X, \mathbb{B}^{n_2+n_3+1}U(1)_{\text{conn}}])_0 \\
\uparrow (1, -1) & & \uparrow \partial \\
\mathbb{Z} & \xrightarrow{\phi_{2,3}} & N([X, \mathbb{B}^{n_2+n_3+1}U(1)_{\text{conn}}])_1
\end{array}
,$$

where the subindices indicate the degree of the chain complex. Now we can represent these chain homotopies succinctly in the upper triangular matrix

$$A = \begin{pmatrix} 0 & \mathcal{G}_1 & \phi_{1,2} & * \\ 0 & 0 & \mathcal{G}_2 & \phi_{2,3} \\ 0 & 0 & 0 & \mathcal{G}_3 \\ 0 & 0 & 0 & 0 \end{pmatrix}.$$

By construction, this matrix satisfies the Maurer-Cartan equation

$$dA - \overline{A} \cdot A = \mu(A) \in \text{Ker}(A).$$

Moreover, $\mu(A)$ is of the form

$$\mu(A) = \begin{pmatrix} 0 & 0 & 0 & \tau \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}.$$

Applying the differential d to τ and using the Leibniz rule, we get

$$\begin{aligned}
d(\tau) &= d(\mathcal{G}_1 \cup \phi_{2,3} - \phi_{1,2} \cup \mathcal{G}_3) \\
&= d(\mathcal{G}_1) \cup \phi_{2,3} + \mathcal{G}_1 \cup d(\phi_{2,3}) - d(\phi_{1,2}) \cup \mathcal{G}_3 + \phi_{1,2} \cup d(\mathcal{G}_3) \\
&= \mathcal{G}_1 \cup (\mathcal{G}_2 \cup \mathcal{G}_3) - (\mathcal{G}_1 \cup \mathcal{G}_2) \cup \mathcal{G}_3 \\
&= 0.
\end{aligned}$$

At the level of sheaf hypercohomology, we have the following:

Proposition 9. *The cohomology class of the matrix cocycle $\mu(A)$ is the element*

$$[\mu(A)] = [\mathcal{G}_1 \cup \phi_{2,3} - \phi_{1,2} \cup \mathcal{G}_3] \in H^{n_1+n_2+n_3+1}(X; U(1)).$$

Proof. We have the following sequence of isomorphisms

$$\begin{aligned}
[\mu(A)] = [\mathcal{G}_1 \cup \phi_{2,3} - \phi_{1,2} \cup \mathcal{G}_3] &\in H_1 \text{hom}_{\mathcal{C}h^+}(C(\mathcal{U}), \mathbb{B}^{n_1+n_2+n_3+2}U(1)_{\text{conn}}) \\
&\simeq \pi_1 \text{Map}(C(\mathcal{U}), \mathbb{B}^{n_1+n_2+n_3+2}U(1)_{\text{conn}}) \\
&\simeq \pi_0 \text{Map}(C(\mathcal{U}), {}^b\mathbb{B}^{n_1+n_2+n_3+1}U(1)) \\
&\simeq H^{n_1+n_2+n_3+1}(X; U(1)).
\end{aligned}$$

□

3.3 General stacky Massey products

We would like to utilize the machinery of May [35] which makes use of matrices. We will introduce stacks labelled by two integers, which will be indexing the entries of the corresponding matrices. To that end, let \mathcal{R}_{ij} , $i, j \in \mathbb{N}$, be simplicial abelian stacks equipped with maps

$$\cup : \mathcal{R}_{ij} \otimes \mathcal{R}_{jk} \rightarrow \mathcal{R}_{ik} ,$$

which are associative in the sense that $\cup \circ (\cup \otimes \text{id}) = \cup \circ (\text{id} \otimes \cup)$.

Remark 7. *Let N denote the normalized Moore functor. It follows from the definition of the differential on the tensor product that the induced product*

$$\tilde{\cup} : N(\mathcal{R}_{ij}) \otimes N(\mathcal{R}_{jk}) \xrightarrow{\sim} N(\mathcal{R}_{ij} \otimes \mathcal{R}_{jk}) \rightarrow N(\mathcal{R}_{ik})$$

must satisfy the Leibniz type rule $d(\alpha \cup \beta) = d(\alpha) \cup \beta + (-1)^{\deg \alpha} \alpha \cup d(\beta)$ on sections.

We can now utilize an extension of the machinery of May [35] locally to define the refined Matric Massey products in our setting. To this end, we consider the set of all upper triangular half-infinite matrices

$$M(\mathcal{R}) = \{A = (a_{ij}), a_{ij} \in N(\mathcal{R}_{ij}), a_{ij} = 0 \text{ for } j \leq i \text{ and } i, j \geq n+1 \text{ for some } n, k \in \mathbb{N}\} . \quad (3.10)$$

Notice that, with our definition, this set possesses more structure. It becomes a sheaf of DGA's with product given by matrix multiplication and differential given by applying the differential on $N(\mathcal{R}_{ij})$ to each entry of the matrix. Just as in the case of classical Massey products, we have a filtration of presheaves of subalgebras

$$M(\mathcal{R})_1 \subset M(\mathcal{R})_2 \subset \dots \subset M(\mathcal{R})_n \subset \dots \quad (3.11)$$

and a bigrading

$$M(\mathcal{R}) = \sum_{p \geq 1, k \geq 0} M^{p,k} , \quad (3.12)$$

where

$$M^{p,k} = \text{span} \left\{ \begin{pmatrix} & & 0 \\ & a_{i,i+p} & \\ 0 & & \end{pmatrix} ; a_{i,i+p} \in N(\mathcal{R}_{i,i+p}) \right\} . \quad (3.13)$$

We can define the following notions similarly to the classical case.

Definition 10. *Let A be a matrix in $M(\mathcal{R})$. We define the (stacky version) of the Maurer-Cartan equation as*

$$dA - \overline{A} \cdot A \equiv 0 \pmod{\ker(A)} ,$$

and call a solution a formal connection with curvature

$$\mu(A) = dA - \overline{A} \cdot A .$$

We are now ready to define the stacky Massey product with a product on the bigraded sequence of stacks.

Definition 11. Let $\mathcal{R} = \{\mathcal{R}_{ij}\}$ be a sequence of abelian stacks equipped with maps

$$\cup : \mathcal{R}_{ij} \otimes \mathcal{R}_{jk} \rightarrow \mathcal{R}_{ik} ,$$

which satisfy $\cup \circ (\text{id} \otimes \cup) = \cup \circ (\cup \otimes \text{id})$. Let A be a formal connection with curvature $\mu(A)$. Then the entries of the hypercohomology class $[\mu(A)]$ are called stacky Massey products.

Remark 8. The following examples of stacks satisfy the compatibility requirement of Def. 11 and will be of particular interest to us. They are the mapping stacks corresponding to the stacks described in Remark 3. Fix a manifold X and a sequence (n_{ij}) , $i < j \leq n$, of integers satisfying $n_{ij} + n_{jk} = n_{ik}$;

1. The stacks $[X, \mathbb{B}^{n_{ij}-1}U(1)_{\text{conn}}]$ of higher bundles with connection, with the stacky cup product and Čech-Deligne differential.
2. The stacks $[X, \mathbb{B}^{n_{ij}}\mathbb{Z}]$ of higher bundles, with the usual cup product and singular differential.
3. The stacks $[X, \flat_{dR}\mathbb{B}^{n_{ij}}U(1)_{\text{conn}}]$ of differential forms of degrees $\leq n$, with the wedge product and exterior derivative.

We highlight the power of the above definitions in the following examples, where we are able to describe all three of the differential, singular, and de Rham triple products.

Example 7. (Differential triple product) Let \mathcal{G}_i , $i = 1, 2, 3$, be bundles $\Delta[0] \rightarrow [X, \mathbb{B}^{n_{i,i+1}-1}U(1)_{\text{conn}}]$. Suppose $\mathcal{G}_1 \cup \mathcal{G}_2$ and $\mathcal{G}_2 \cup \mathcal{G}_3$ represent trivial classes in $\pi_0\text{Map}(X, \mathbb{B}^{n_{ij}-1}U(1)_{\text{conn}})$. Choose a defining system

$$A = \begin{pmatrix} 0 & \mathcal{G}_1 & \phi_{1,2} & * \\ 0 & 0 & \mathcal{G}_2 & \phi_{2,3} \\ 0 & 0 & 0 & \mathcal{G}_3 \\ 0 & 0 & 0 & 0 \end{pmatrix} ,$$

where $\phi_{1,2}$ and $\phi_{2,3}$ are nondegenerate 1-simplices trivializing the cup products. Then the curvature of A is

$$\mu(A) = \begin{pmatrix} 0 & 0 & 0 & \tau \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix} ,$$

and the hypercohomology class $[\tau]$ is $[\mathcal{G}_1 \cup \phi_{2,3} - \phi_{1,2} \cup \mathcal{G}_3]$. The latter is an element in

$$\pi_1\text{Map}(X, \mathbb{B}^{n_{33}-1}U(1)_{\text{conn}}) \simeq \pi_0\text{Map}(X, \flat\mathbb{B}^{n_{14}-2}U(1)) \simeq H^{n_{14}-2}(X; U(1)) ,$$

where we have $n_{14} = n_{13} + n_{34} = n_{12} + n_{23} + n_{34}$.

Example 8. (Singular triple product) Let X be a manifold, and let $|X|$ be the topological space denoting its geometric realization. Let $a_i : |X| \rightarrow K(\mathbb{Z}, n_{i,i+1}) \simeq B^{n_{i,i+1}}\mathbb{Z}$, $i = 1, 2, 3$, be singular cochains with cup products vanishing in cohomology. Choose a defining system

$$A = \begin{pmatrix} 0 & a_1 & f_{1,2} & * \\ 0 & 0 & a_2 & f_{2,3} \\ 0 & 0 & 0 & a_3 \\ 0 & 0 & 0 & 0 \end{pmatrix} .$$

Since geometric realization is a left ∞ -adjoint the discrete stack functor disc [47], these are equivalently given by maps of stacks

$$\bar{a}_i : \Delta[0] \rightarrow [X, \mathbb{B}^{n_{i,i+1}} \mathbb{Z}] ,$$

and homotopies

$$\bar{f}_{i,i+1} : \Delta[1] \rightarrow [X, \mathbb{B}^{n_{i,i+2}} \mathbb{Z}]$$

trivializing the cup products, hence a defining system

$$A = \begin{pmatrix} 0 & \bar{a}_1 & \bar{f}_{1,2} & * \\ 0 & 0 & \bar{a}_2 & \bar{f}_{2,3} \\ 0 & 0 & 0 & \bar{a}_3 \\ 0 & 0 & 0 & 0 \end{pmatrix} .$$

The hypercohomology class of the entry $\tau \in \mu(A)$ is given by $[\bar{a}_1 \cup \bar{f}_{2,3} - \bar{f}_{1,2} \cup \bar{a}_3]$, which is an element in

$$\begin{aligned} \pi_1 \text{Map}(X, \mathbb{B}^{n_{14}} \mathbb{Z}) &\simeq \pi_1 \text{Map}(|X|, K(\mathbb{Z}, n_{14})) \\ &\simeq \pi_0 \text{Map}(|X|, K(\mathbb{Z}, n_{14} - 1)) \\ &\simeq H^{n_{14}-1}(X, \mathbb{Z}) . \end{aligned}$$

Example 9. (de Rham triple product) Let X be a manifold and let α_i , $i = 1, 2, 3$, be closed forms in different degrees. These forms are equivalently given by maps

$$\alpha_i : \Delta[0] \rightarrow \mathfrak{b}_{dR} \mathbb{B}^{n_{i,i+1}} U(1)_{\text{conn}} .$$

Suppose that the wedge products $\alpha_1 \wedge \alpha_2$ and $\alpha_2 \wedge \alpha_3$ are trivial in cohomology. Then we can choose a defining system via

$$A = \begin{pmatrix} 0 & \alpha_1 & \eta_{1,2} & * \\ 0 & 0 & \alpha_2 & \eta_{2,3} \\ 0 & 0 & 0 & \alpha_3 \\ 0 & 0 & 0 & 0 \end{pmatrix} ,$$

where $\eta_{1,2}$ and $\eta_{2,3}$ are 1-simplices. The hypercohomology class of the entry $\tau \in \mu(A)$ is given by

$$[\alpha_1 \wedge \eta_{2,3} - \eta_{1,2} \cup \alpha_3] .$$

Since the sheaf $\Omega^{\leq n_{14}}$ is fine in each degree, we can calculate the hypercohomology as

$$\begin{aligned} \pi_1 \text{Map}(X, \Omega^{\leq n_{14}}) &\simeq H_1 \Omega^{\leq n_{14}}(X) \\ &\simeq H_{dR}^{n_{14}-1}(X) . \end{aligned}$$

Our main result in this section relates Massey products for Deligne cocycles to corresponding ones for higher bundles in the stacky sense.

Theorem 12. Let \hat{a}_i , $1 \leq i \leq l$, be Deligne cocycles. Suppose the l -fold Massey product is defined. Let \mathcal{G}_i , $1 \leq i \leq l$, be $n_{i,i+1}$ -bundles with connections

$$\mathcal{G}_i : X \rightarrow \mathbb{B}^{n_{i,i+1}} U(1)_{\text{conn}} ,$$

representing the Deligne cocycles. Then there is a natural bijection between corresponding Massey products

$$\langle \mathcal{G}_1, \mathcal{G}_2, \dots, \mathcal{G}_l \rangle \simeq \langle \hat{a}_1, \hat{a}_2, \dots, \hat{a}_l \rangle .$$

Proof. Recall that $\mathbb{B}^n U(1)_{\text{conn}} := \Gamma(\mathbb{Z}_{\mathcal{D}}^\infty[n+1])$ (see [21]). Using the definition of the stacky hom, the fact that the counit $\epsilon : N\Gamma \rightarrow \text{id}$ is a natural isomorphism and the lax monoidal structure on N , we have a homotopy equivalence for each test object U ,

$$\begin{aligned} N([X, \mathbb{B}^n U(1)_{\text{conn}}])(U) &= N([X, \mathbb{B}^n U(1)_{\text{conn}}](U)) \\ &\simeq N(\text{Map}(C(\{U_i\}) \times U, \mathbb{B}^n U(1)_{\text{conn}})) \\ &\simeq \text{hom}_{\text{ch}^+}(N(C(\{U_i\}) \otimes N(U), \mathbb{Z}_{\mathcal{D}}^\infty[n+1])) \\ &\simeq \text{hom}_{\text{ch}^+}(N(C(\{U_i\}), \mathbb{Z}_{\mathcal{D}}^\infty[n+1])(U)) \\ &\simeq C(X, \mathbb{Z}_{\mathcal{D}}^\infty[n+1])(U) , \end{aligned}$$

where the last line denotes the Čech resolution of the Deligne complex $\mathbb{Z}_{\mathcal{D}}^\infty[n+1]$. Hence, a defining system in the stacky sense is naturally equivalent to a defining system in the sense of [44]. Since the set of Massey products is parametrized by the set of defining systems, it follows that indeed we have a natural bijection

$$\langle \mathcal{G}_1, \mathcal{G}_2, \dots, \mathcal{G}_l \rangle \simeq \langle \hat{a}_1, \hat{a}_2, \dots, \hat{a}_l \rangle .$$

□

3.4 Properties of stacky Massey products

We will now consider properties of the stacky Massey products. Our setting allows for these to be quite attractive and natural. The most immediate of those are direct generalizations of classical ones. Later in this section we will see properties that are more peculiar to the differential setting. Among the properties that the classical Massey products satisfy are the following (see [35] [30]):

1. *Dimension:* The dimension of $\langle x_1, x_2, \dots, x_n \rangle$ is $\sum \deg(x_i) - n + 2$.
2. *Naturality:* If $f : X \rightarrow Y$ is a continuous map and $y_1, \dots, y_k \in H^*(Y; R)$ such that the k -fold Massey product $\langle y_1, y_2, \dots, y_k \rangle$ is defined, then $\langle x_1, \dots, x_k \rangle = \langle f^*(y_1), \dots, f^*(y_k) \rangle$ is defined as a Massey product on the cohomology of X and

$$f^*(\langle y_1, \dots, y_k \rangle) \subset \langle f^*(y_1), \dots, f^*(y_k) \rangle .$$

3. *Definedness:* The vanishing of the the lower Massey products is only a necessary condition for the k -fold Massey product to be defined for $k > 3$. For $k = 3$ the condition is both necessary and sufficient.
4. *Slide relation:* If the Massey product $\langle x_1, x_2, \dots, x_n \rangle$ is defined, then so is $\langle x_1, x_2, \dots, rx_i, \dots, x_n \rangle$ for any $r \in R$. Moreover we have the relation

$$r\langle x_1, x_2, \dots, x_n \rangle \subset \langle x_1, x_2, \dots, rx_i, \dots, x_n \rangle .$$

These indeed extends to the stacky version.

Proposition 13. *The stacky Massey products satisfy the following properties:*

1. *Dimension:* The dimension of $\langle \mathcal{G}_1, \mathcal{G}_2, \dots, \mathcal{G}_l \rangle$ is $\sum \deg(\mathcal{G}_i) - l + 2$.

2. Naturality: If $f : X \rightarrow Y$ is a smooth map between manifolds and $\mathcal{G}_1, \dots, \mathcal{G}_k \in \hat{H}_{\mathcal{D}}^*(X; \mathbb{Z})$ such that the k -fold Massey product $\langle \mathcal{G}_1, \mathcal{G}_2, \dots, \mathcal{G}_k \rangle$ is defined, then $\langle \mathcal{G}_1, \dots, \mathcal{G}_k \rangle = \langle f^*(\mathcal{G}_1), \dots, f^*(\mathcal{G}_k) \rangle$ is defined as a Massey product on the differential cohomology of X and

$$f^*(\langle \mathcal{G}_1, \dots, \mathcal{G}_k \rangle) \subset \langle f^*(\mathcal{G}_1), \dots, f^*(\mathcal{G}_k) \rangle .$$

3. Definedness: The vanishing of the lower Massey products is only a necessary condition for the k -fold Massey product to be defined for $k > 3$. For $k = 3$ the condition is both necessary and sufficient.
4. Slide relation: If the Massey product $\langle \mathcal{G}_1, \mathcal{G}_2, \dots, \mathcal{G}_n \rangle$ is defined, then so is $\langle \mathcal{G}_1, \mathcal{G}_2, \dots, m\mathcal{G}_i, \dots, \mathcal{G}_n \rangle$ for any $m \in \mathbb{Z}$. Moreover we have the relation

$$m\langle \mathcal{G}_1, \mathcal{G}_2, \dots, \mathcal{G}_n \rangle \subset \langle \mathcal{G}_1, \mathcal{G}_2, \dots, m\mathcal{G}_i, \dots, \mathcal{G}_n \rangle .$$

Proof. Part 1 follows immediately from the definition. To prove part 2, note that the functor $[-, \mathcal{R}]$ is contravariant, sending a map $f : X \rightarrow Y$ to its pullback

$$f^* : [Y, \mathcal{R}_{ij}] \rightarrow [X, \mathcal{R}_{ij}] .$$

Since the cup product is natural with respect to pullbacks, the induced morphism $f^* : N([Y, \mathcal{R}_{ij}]) \rightarrow N([X, \mathcal{R}_{ij}])$ descends to a morphism of sheaves of DGA's

$$f^* : M([Y, \mathcal{R}_{ij}]) \rightarrow M([X, \mathcal{R}_{ij}]) .$$

It follows that if A is a formal connection in $M([Y, \mathcal{R}_{ij}])$, then $f^*(A)$ is a formal connection in $M([X, \mathcal{R}_{ij}])$ satisfying the equation:

$$df^*(A) - f^*(A) \cdot \overline{f^*(A)} = f^*(\mu(A)) \in \ker(f^*(A)) .$$

By definition of the k -fold Massey product, the claim follows. For part 3, we will show that for $k = 3$ the condition is both necessary and sufficient. From the proof, it will be clear that this cannot be the case for higher products. Let $\mathcal{G}_1, \mathcal{G}_2$ and \mathcal{G}_3 be bundles and suppose the triple product $\langle \mathcal{G}_1, \mathcal{G}_2, \mathcal{G}_3 \rangle$ is defined. Then we have trivializations $\phi_{1,2}$ and $\phi_{2,3}$ such that

$$d\phi_{1,2} = \mathcal{G}_1 \cup \mathcal{G}_2 , \quad d\phi_{2,3} = \mathcal{G}_2 \cup \mathcal{G}_3 . \quad (3.14)$$

Hence, both cup products are trivial. For the converse, it is clear that if both cup products are trivial in cohomology, we can choose trivializing homotopies and form the Massey triple product. For higher products, the higher trivializations depend on the lower ones. In fact, for the fourfold product, choose trivializations $\phi_{1,2}$, $\phi_{2,3}$ and $\phi_{3,4}$ of the cup products such that

$$\mathcal{G}_1 \cup \phi_{2,3} - \phi_{1,2} \cup \mathcal{G}_3 \in \langle \mathcal{G}_1, \mathcal{G}_2, \mathcal{G}_3 \rangle$$

is trivializable. Then for the fourfold Massey product to be defined, the other triple product

$$\mathcal{G}_2 \cup \phi_{2,3} - \phi_{3,4} \cup \mathcal{G}_4 \in \langle \mathcal{G}_2, \mathcal{G}_3, \mathcal{G}_4 \rangle$$

must be trivializable. But this may not be true, even if $\langle \mathcal{G}_2, \mathcal{G}_3, \mathcal{G}_4 \rangle$ contains 0. Finally, for part 4, let A be a formal connection of the form

$$A = \begin{pmatrix} 0 & \mathcal{G}_1 & \phi_{1,2} & \dots & & & & & * \\ & 0 & \mathcal{G}_2 & \phi_{2,3} & \dots & & & & \\ & & 0 & \dots & \dots & \dots & & & \\ & & & & \mathcal{G}_{i-1} & \phi_{i-1,i} & & & \\ & & & & & \mathcal{G}_i & \phi_{i,i+1} & & \\ & & & & & & \mathcal{G}_{i+1} & & \\ & & & & & & & \dots & \phi_{n-2,n-1} \\ & & & & & & & & \mathcal{G}_{n-1} & \phi_{n-1,n} \\ & & & & & & & & & \mathcal{G}_n \\ & & & & & & & & & 0 \end{pmatrix}.$$

Then the matrix

$$\tilde{A} = \left(\begin{array}{cccccc|cccccc} 0 & \mathcal{G}_1 & \phi_{1,2} & \dots & \dots & m\phi_{1,i} & \dots & & & & * \\ & 0 & \mathcal{G}_2 & \phi_{2,3} & \dots & \dots & & & & & \\ & & 0 & \dots & \dots & m\phi_{i-2,i} & \dots & & & & \\ & & & & \mathcal{G}_{i-1} & m\phi_{i-1,i} & m\phi_{i-1,i+1} & \dots & & & \\ & & & & & m\mathcal{G}_i & m\phi_{i,i+1} & m\phi_{i,i+2} & \dots & m\phi_{i,n} & \\ \hline & & & & & & \mathcal{G}_{i+1} & \dots & & & \\ & & & & & & & \dots & \phi_{n-2,n-1} & & \\ & & & & & & & 0 & \mathcal{G}_{n-1} & \phi_{n-1,n} & \\ & & & & & & & & 0 & \mathcal{G}_n & \\ & & & & & & & & & 0 & \end{array} \right)$$

is also a formal connection: that is, a defining system for the Massey product $\langle \mathcal{G}_1, \dots, m\mathcal{G}_i, \dots, \mathcal{G}_n \rangle$. Indeed, let us write the matrix A as a block matrix

$$A = \left(\begin{array}{c|c} A_1 & A_2 \\ \hline 0 & A_3 \end{array} \right).$$

Then the second matrix can be written

$$\tilde{A} = \left(\begin{array}{c|c} A_1 & mA_2 \\ \hline 0 & A_3 \end{array} \right).$$

Now the Maurer-Cartan equation for \tilde{A} reads

$$\begin{aligned}\mu(\tilde{A}) &= \left(\begin{array}{c|c} dA_1 & mdA_2 \\ \hline 0 & dA_3 \end{array} \right) - \left(\begin{array}{c|c} \overline{A_1} & \overline{mA_2} \\ \hline 0 & \overline{A_3} \end{array} \right) \left(\begin{array}{c|c} A_1 & mA_2 \\ \hline 0 & A_3 \end{array} \right) \\ &= \left(\begin{array}{c|c} dA_1 & mdA_2 \\ \hline 0 & dA_3 \end{array} \right) - \left(\begin{array}{c|c} \overline{A_1}A_1 & m(\overline{A_1}A_2 + \overline{A_2}A_3) \\ \hline 0 & \overline{A_3}A_3 \end{array} \right).\end{aligned}$$

We want to show that $\mu(\tilde{A})$ is in $\ker(\tilde{A})$. Since A satisfies the Maurer-Cartan equation up to an element in the kernel

$$\ker(A) = \begin{pmatrix} \dots & 0 & * \\ \dots & 0 & 0 \\ & \vdots & \vdots \end{pmatrix} \quad (3.15)$$

we must have $dA_1 = \overline{A_1} \cdot A_1$ and $dA_3 = \overline{A_3} \cdot A_3$. Since A is formal, we must also have

$$\mu(A)_2 = dA_2 - (\overline{A_1}A_2 + \overline{A_2}A_3)$$

where $\mu(A)_2$ is the upper right block of $\mu(A)$ of dimension $\dim(A_2)$. Since the only nonzero term of $\mu(A)$ is the cochain representative of the Massey product τ , located in the upper right corner of $\mu(A)$, we have that

$$\mu(\tilde{A})_2 = mdA_2 - m(A_1\overline{A_2} + A_2\overline{A_3}) = m\mu(A)_2$$

has one nonzero element $\sigma = m\tau$ in the upper right corner. Therefore, \tilde{A} is indeed formal and, at the level of cohomology, the only nonzero term of the class $[\mu(A)]$ is $[\sigma] = m[\tau]$. Since $[\tau]$ was chosen to be an arbitrary element of the Massey product $\langle \mathcal{G}_1, \dots, \mathcal{G}_n \rangle$, we have

$$m\langle \mathcal{G}_1, \dots, \mathcal{G}_n \rangle \subset \langle \mathcal{G}_1, \dots, m\mathcal{G}_i, \dots, \mathcal{G}_n \rangle.$$

□

We now discuss the relationship between the stacky Massey product and the singular Massey product. The following parametrizes how forgetting the differential data on the Massey product is not quite the same as taking the Massey product of cohomology classes after forgetting the differential data on these.

Proposition 14. *Let $\mathcal{G}_i : \Delta[0] \rightarrow [X, \mathbb{B}^{n_{i,i+1}}U(1)_{\text{conn}}]$, $1 \leq i \leq l$, be higher bundles on X with defined Massey product. Then precomposition with the forgetful morphism*

$$I : \mathbb{B}^n U(1)_{\text{conn}} \rightarrow \mathbb{B}^{n+1} \mathbb{Z},$$

induced by the map

$$\begin{array}{ccc}
\underline{\mathbb{Z}} & \xrightarrow{0} & \underline{\mathbb{Z}} \\
\downarrow i & & \downarrow d \\
\Omega^0 & \xrightarrow{0} & 0 \\
\downarrow & & \downarrow \\
\vdots & \xrightarrow{0} & \vdots \\
\downarrow & & \downarrow \\
\Omega^{n-1} & \xrightarrow{d} & 0,
\end{array}$$

yields singular cocycles with defined Massey product. Furthermore, we have

$$I\langle \mathcal{G}_1, \mathcal{G}_2, \mathcal{G}_3 \rangle \subset \langle I(\mathcal{G}_1), I(\mathcal{G}_2), I(\mathcal{G}_3) \rangle.$$

Proof. For simplicity, we denote the sheaf of matrix algebras

$$\begin{aligned}
M_{\text{diff}} &:= M([X, \mathbb{B}^*U(1)_{\text{conn}}]) \\
M_{\text{sing}} &:= M([X, \mathbb{B}^*\mathbb{Z}])
\end{aligned}$$

according to the corresponding cohomology theories for these matrices. It is clear by definition that I respects the cup product structure, hence I induces a morphism of sheaves of DGA's $I_* : M_{\text{diff}} \rightarrow M_{\text{sing}}$. It follows immediately from the definition of the Maurer-Cartan equation Def. 10, that formal connections are sent to formal connections. Then passing to hypercohomology gives the result. \square

It follows from the proposition that if the classical Massey product $\langle I(\mathcal{G}_1), I(\mathcal{G}_2), I(\mathcal{G}_3) \rangle$ is zero then certainly the left hand side is zero, i.e. $\langle \mathcal{G}_1, \mathcal{G}_2, \mathcal{G}_3 \rangle$ is in the kernel of the forgetful morphism I . From the sequence $\Omega^{n-1}/\text{Im}(d) \rightarrow \hat{H}^n \xrightarrow{I} H^n$ we have that $\langle \mathcal{G}_1, \mathcal{G}_2, \mathcal{G}_3 \rangle$ will be an $(n-1)$ -form. However, it is important to note that this is not quite the $(n-1)$ -form given by the classical Massey product. A related question is to ask whether the differential Massey product completely refines the singular Massey product. That is: do we have a *bijection*,

$$I\langle \mathcal{G}_1, \mathcal{G}_2, \mathcal{G}_3 \rangle \simeq \langle I(\mathcal{G}_1), I(\mathcal{G}_2), I(\mathcal{G}_3) \rangle ?$$

Unfortunately, this cannot be possible. Essentially, this is because the map $I_* : M_{\text{diff}} \rightarrow M_{\text{sing}}$ has a nontrivial kernel. Hence we cannot expect the Maurer-Cartan equation to hold after refining. However, this does help explain the nature of differential Massey products. In fact, since these products are always flat, it follows from diagram (2.7) that *if* the refinement of a singular formal connection is again a formal connection, then the singular Massey product must have been torsion. We will show that the failure of the refinement to satisfy the Maurer-Cartan equation can be measured by the de Rham Massey product.

Lemma 15. *Let $\mathcal{F}_{ij} \rightarrow \mathcal{R}_{ij} \twoheadrightarrow \mathcal{S}_{ij}$ be a fibration sequence of abelian prestacks for each i and j . Suppose, moreover, that we have commuting diagrams*

$$\begin{array}{ccc}
\mathcal{F}_{ij} \otimes \mathcal{F}_{jk} & \xrightarrow{\cup} & \mathcal{F}_{ik} \\
\downarrow i \otimes i & & \downarrow i \\
\mathcal{R}_{ij} \otimes \mathcal{R}_{jk} & \xrightarrow{\cup} & \mathcal{R}_{ik} \\
\downarrow p \otimes p & & \downarrow p \\
\mathcal{S}_{ij} \otimes \mathcal{S}_{jk} & \xrightarrow{\cup} & \mathcal{S}_{ik} .
\end{array}$$

Then the induced sequence $0 \rightarrow M(\mathcal{F}) \rightarrow M(\mathcal{R}) \twoheadrightarrow M(\mathcal{S}) \rightarrow 0$ is a short exact sequence of DGA's.

Proof. Since the normalized Moore functor is right Quillen and preserves equivalences, it follows that it sends fiber sequences to fiber sequences. Hence, we have a diagram

$$\begin{array}{ccc}
N(\mathcal{F}_{ij}) \otimes N(\mathcal{F}_{jk}) & \xrightarrow{\cup} & N(\mathcal{F}_{ik}) \\
\downarrow i \otimes i & & \downarrow i \\
N(\mathcal{R}_{ij}) \otimes N(\mathcal{R}_{jk}) & \xrightarrow{\cup} & N(\mathcal{R}_{ik}) \\
\downarrow p \otimes p & & \downarrow p \\
N(\mathcal{S}_{ij}) \otimes N(\mathcal{S}_{jk}) & \xrightarrow{\cup} & N(\mathcal{S}_{ik}) ,
\end{array}$$

where the right hand side is a short exact sequence of presheaves of chain complexes. By definition, it follows that we have a short exact sequence

$$0 \rightarrow M(\mathcal{F}) \rightarrow M(\mathcal{R}) \twoheadrightarrow M(\mathcal{S}) \rightarrow 0$$

of chain complexes. By commutivity of the above diagram, both maps are homomorphisms of presheaves of DGA's. \square

It follows from the lemma along with diagram (2.19), that there is a short exact sequence of presheaves of bigraded rings

$$0 \rightarrow M([X, \Omega^{\leq*}]) \rightarrow M_{\text{diff}} \rightarrow M_{\text{sing}} \rightarrow 0 . \quad (3.16)$$

Hence, $M_{\text{form}} := M([X, \Omega^{\leq*}])$ is a two-sided ideal in M_{diff} .

Now, by definition of $\ker(A)$ along with the above observation, we have

$$\ker(\hat{A}) \subset \widehat{\ker(A)} ,$$

where $\widehat{}$ denotes a choice of differential refinement. In fact, for a matrix $C \in \ker(\hat{A})$ and $C' \in \widehat{\ker(\hat{A})}$, we have that the difference $C - C' = B \in M_{\text{form}}$. It is this lack of commutativity between taking kernels and taking differential refinements that leads to a nontrivial structure than might otherwise be anticipated.

Summarizing the previous observations gives the following theorem.

Theorem 16. *Let A be a formal connection for M_{sing} , and let \hat{A} be a differential refinement of A with $\mu(A)$ a solution to the Maurer-Cartan equation. Then any differential refinement $\widehat{\mu(A)}$ satisfies the twisted Maurer-Cartan equation*

$$\widehat{\mu(A)} = d\hat{A} - \overline{\hat{A}} \cdot \hat{A} \equiv B \pmod{\ker(\hat{A})}, \quad (3.17)$$

where B is some matrix in the ideal M_{form} .

Proof. Since A is formal, $\mu(A)$ satisfies

$$\mu(A) = dA - \overline{A} \cdot A \equiv 0 \pmod{\ker(A)}.$$

Hence, any refinement must satisfy

$$\widehat{\mu(A)} = D\hat{A} - \overline{\hat{A}} \cdot \hat{A} \equiv 0 \pmod{\widehat{\ker(A)}},$$

where $D = d + (-1)^*\delta$ is the Čech-Deligne differential on M_{diff} . Now by sequence (3.16), we see that this is equivalent to existence of a matrix of forms B satisfying (3.17). \square

In general, the Deligne-Beilinson cup product does not refine the de Rham wedge product for the whole de Rham complex, but does so only for the top and bottom degrees, as we have seen in Prop. 5 and Prop. 6. However, for the triple product the only cup products that arise are between degree zero and degree one cocycles, so that nothing is missed in passing to \cup_{DB} . Consequently, for the case of the triple product, the matrix B in the above example encodes the information needed to define the de Rham Massey product. More precisely, we have the following.

Proposition 17. *Let $a_i \in H^*(X, \mathbb{Z})$, $i = 1, 2, 3$, and let $\iota(a)_i \in H_{\text{dR}}^*(X)$ denote the inclusions into de Rham cohomology. Let*

$$A = \begin{pmatrix} 0 & a_1 & \phi_{1,2} & * \\ 0 & 0 & a_2 & \phi_{2,3} \\ 0 & 0 & 0 & a_3 \\ 0 & 0 & 0 & 0 \end{pmatrix}.$$

in M_{sing} be a matrix of singular cochains defining a formal connection and let $\mu(A)$ be the corresponding solution to the corresponding Maurer-Cartan equation. Then for any differential refinement $\widehat{\mu(A)}$ of $\mu(A)$, the curvature $R(\widehat{\mu(A)})$ is a de Rham Massey product in $\langle \iota(a)_1, \iota(a)_2, \iota(a)_3 \rangle$. If, in addition, $\widehat{\mu(A)}$ is a solution to the differential Maurer-Cartan equation, then $R(\widehat{\mu(A)}) = 0$ and $\mu(A)$ represents a torsion class.

Proof. Let a_i , $i = 1, 2, 3$, be singular cochains of degree $n_{i,i+1}$. Suppose that the triple product is defined, and choose a defining system

$$A = \begin{pmatrix} 0 & a_1 & \phi_{1,2} & * \\ 0 & 0 & a_2 & \phi_{2,3} \\ 0 & 0 & 0 & a_3 \\ 0 & 0 & 0 & 0 \end{pmatrix}.$$

Let

$$\hat{A} = \begin{pmatrix} 0 & \hat{a}_1 & \hat{\phi}_{1,2} & * \\ 0 & 0 & \hat{a}_2 & \hat{\phi}_{2,3} \\ 0 & 0 & 0 & \hat{a}_3 \\ 0 & 0 & 0 & 0 \end{pmatrix}$$

be a refinement. Then we know that the refinement $\widehat{\mu(A)}$ satisfies the equation $D\hat{A} = \hat{A} \cdot \overline{\hat{A}} + B$ up to some element in $\ker(\hat{A})$. Explicitly, letting $B = (\eta_{ij})$, we have

$$\hat{A} = \begin{pmatrix} 0 & 0 & D\hat{\phi}_{1,2} & * \\ 0 & 0 & 0 & D\hat{\phi}_{2,3} \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix} = \begin{pmatrix} 0 & \eta_{12} & \hat{a}_1 \cup \hat{a}_2 + \hat{\eta}_{13} & \hat{a}_1 \cup \hat{\phi}_{2,3} - \hat{\phi}_{1,2} \cup \hat{a}_3 \\ 0 & 0 & \eta_{23} & \hat{a}_1 \cup \hat{a}_2 + \hat{\eta}_{24} \\ 0 & 0 & 0 & \eta_{34} \\ 0 & 0 & 0 & 0 \end{pmatrix}.$$

The requirement that this equation hold up to an element in $\ker(\hat{A})$ forces the equations

$$\begin{aligned} \eta_{12} &= 0, & D\hat{\phi}_{1,2} &= \hat{a}_1 \cup \hat{a}_2 + \eta_{13}, \\ \eta_{23} &= 0, & D\hat{\phi}_{2,3} &= \hat{a}_1 \cup \hat{a}_2 + \eta_{24}. \end{aligned}$$

At the level of connections, the data provided by the right two equations reduces to

$$d\phi_{1,2} = b_1 \wedge a_2 + \eta_{13} \tag{3.18}$$

$$d\phi_{2,3} = b_2 \wedge a_3 + \eta_{24}, \tag{3.19}$$

where b_1 and b_2 are forms representing the connections with curvatures a_1 and a_2 .

Now forming $\widehat{\mu(A)}$ gives the matrix

$$\widehat{\mu(A)} = \begin{pmatrix} 0 & 0 & 0 & \hat{a}_1 \cup \hat{\phi}_{2,3} - \hat{\phi}_{1,2} \cup \hat{a}_3 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}.$$

Finally, applying the curvature map R to the only nonzero term gives

$$\begin{aligned} R(\hat{a}_1 \cup \hat{\phi}_{2,3} - \hat{\phi}_{1,2} \cup \hat{a}_3) &= R(\hat{a}_1) \wedge R(\hat{\phi}_{2,3}) - R(\hat{\phi}_{1,2}) \wedge R(\hat{a}_3) \\ &= R(\hat{a}_1) \wedge R(\hat{a}_2 \cup \hat{a}_3) + R(\hat{a}_1) \wedge \eta_{24} - (\eta_{13} \wedge R(\hat{a}_3) + R(\hat{a}_1 \cup \hat{a}_2) \wedge R(\hat{a}_3)) \\ &= R(\hat{a}_1 \cup (\hat{a}_2 \cup \hat{a}_3)) + R(\hat{a}_1) \wedge \eta_{24} - \eta_{13} \wedge R(\hat{a}_3) - R((\hat{a}_1 \cup \hat{a}_2) \cup \hat{a}_3) \\ &= R(\hat{a}_1) \wedge \eta_{24} - \eta_{13} \wedge R(\hat{a}_3) \\ &= a_1 \wedge \eta_{24} - \eta_{13} \wedge a_3. \end{aligned}$$

Notice that it follows from equations (3.18) and (3.19) that the last line represents a de Rham Massey product (simply apply d to both sides of those equations). This proves the first claim.

For the second, observe that if $\hat{\mu}$ solves the Maurer-Cartan equation, then we can choose $B = (\eta_{ij}) = 0$, and the curvature calculated above must vanish. \square

4 Applications

We will discuss our applications in this section, both from geometry and mathematical physics. We will show how Massey products arise in various settings, both classically and then in the newly constructed stacky form.

4.1 Trivializations for (higher) structures

In this section we will consider Massey products arising from characteristic classes, hence associated with bundles or (higher) abelian gerbes. The refined Massey products will be associated with bundles or (higher) abelian gerbes together with connections on them.

We begin with a toy example.

Example 10. *[Circle bundles] Let $S^1 \rightarrow E \xrightarrow{\pi} \Sigma$ be a circle bundle over an orientable closed surface Σ of genus $g > 0$. Suppose that the first Chern class of E is nonzero. The Gysin sequence implies that $H^1(\Sigma, \mathbb{R})$ is isomorphic to $H^1(E, \mathbb{R})$ and that $H^2(\Sigma, \mathbb{R})$ maps to zero in $H^2(E, \mathbb{R})$. Let $a, b \in H^1(\Sigma, \mathbb{R})$ such that $\pi^*c_1(E) = a \cup b$. Then by Chern-Weil theory there is a connection on E whose 1-form A satisfies $dA = \pi^*c_1(E) = a \cup b$. Moreover, since the Gysin sequence holds for both singular and de Rham cohomology, $a \cup b$ vanishes in both singular and de Rham cohomology and we can assume that A represents an integral cochain. Then we consider the pairing*

$$(A \cup a \cup b)([E]) = (A \cup dA)[E] = c_1(E)([\Sigma]) \neq 0.$$

Hence we obtain the Massey product $\langle a, a, b \rangle$ which is represented by $A \cup a$.

We now discuss the differential refinement of the above example. This will involve the Deligne derivative D , which in the setting of the Čech-de Rham double complex, is given by $D = d + (-1)^k \delta$.

Example 11. *Let \hat{a} and \hat{b} be differential refinements of a and b . Suppose, moreover, that we have*

$$\pi^*\hat{c}_1(E) = \hat{a} \cup \hat{b} \quad \text{and} \quad D\hat{A} = \pi^*\hat{c}_1(E) = \hat{a} \cup \hat{b} \tag{4.1}$$

so that $\pi^*\hat{c}_1(E)$ is trivializable as a bundle equipped with connection. Note that in the context of the previous example the first equation is true, at least in the case where \hat{a} and \hat{b} arise as characteristic forms (see [6]). However, the second equation is much stronger than the requirement that $\pi^*c_1(E)$ vanish in usual cohomology. For example, this equation implies that the bundle $\pi^*\hat{c}_1(E)$ (with curvature $\pi^*c_1(E)$) is flat. We also make one additional assumption which may seem, a priori, unrelated but is in fact necessary to define Massey products of this type. We assume that \bar{a} is annihilated by the first Steenrod square $Sq^1(\bar{a}) = 0$, that is \bar{a} is the mod 2 reduction of an integral class a . It was shown by Gomi [25] that for a differential cohomology classes \hat{a} of odd degree n , we have the formula

$$\hat{a} \cup \hat{a} = i(Sq^{n-1}(\bar{a})).$$

Here, the right hand side is to be understood as the inclusion of $Sq^{n-1}(\bar{a})$ into $U(1)$ cohomology via the obvious morphism $\mathbb{Z}/2 \rightarrow U(1)$ (see [25][6]), and i denotes the inclusion into differential cohomology via the map in diagram (2.7) (which raises the degree by 1).

Now let ϕ be a trivialization of $Sq^1(\bar{a})$. Since the inclusion into differential cohomology holds at the level of chains and is natural, we have in our degrees

$$d(i(\phi)) = id(\phi) = i(Sq^1(\bar{a})) = \hat{a} \cup \hat{a}. \tag{4.2}$$

Now the following matrix organizes the defining system given by equations (4.1) and (4.2):

$$\begin{pmatrix} 0 & \hat{a} & i(\phi) & \\ 0 & 0 & \hat{a} & \hat{A} \\ 0 & 0 & 0 & \hat{b} \\ 0 & 0 & 0 & 0 \end{pmatrix}.$$

Then an element of the Massey product $\langle \hat{a}, \hat{a}, \hat{b} \rangle$ is given by the class

$$\hat{a} \cup \hat{A} - i(\phi) \cup \hat{b}.$$

Both of the previous examples can be generalized to higher Chern classes.

Example 12. Let n be an odd integer and let $E \xrightarrow{\pi} \Sigma_{2n}$ be a complex vector bundle over an orientable closed $2n$ manifold Σ_{2n} . Suppose that the n -th Chern class of the E is nonzero. Let $a_n, b_n \in H^n(\Sigma_{2n}, \mathbb{R})$ be such that $\pi^*c_n(E) = a \cup b$. Then there is a $(2n-1)$ -form $A_{(2n-1)}$ representing a singular cochain such that $dA_{(2n-1)} = \pi^*c_n(E) = a_n \cup b_n$. Then we consider the pairing

$$(A_{(2n-1)} \cup a_n \cup b_n)([E]) = (A_{(2n-1)} \cup dA_{(2n-1)})([E]) = c_n(E)([\Sigma]) \neq 0.$$

Hence we obtain the Massey product $\langle a, a, b \rangle$ which is represented by $A_{(2n-1)} \cup a_n$.

Example 13. Let \hat{a}_n and \hat{b}_n be differential refinements of a_n and b_n . Suppose moreover that we have

$$\pi^*\hat{c}_n(E) = \hat{a}_n \cup \hat{b}_n \text{ and } d\hat{A}_{(2n-1)} = \pi^*\hat{c}_n(E) = \hat{a}_n \cup \hat{b}_n, \quad (4.3)$$

so that $\pi^*\hat{c}_n(E)$ is trivializable as a bundle equipped with connection. We also assume that $Sq^{n-1}(\bar{a}_n) = 0$, where \bar{a} is the mod 2 reduction of a . Then we again have $\hat{a}_n \cup \hat{a}_n = i(Sq^{n-1}(\bar{a}_n))$, by Gomi [25]. Now let ϕ be a trivialization of $Sq^{n-1}(\bar{a}_n)$. Since the inclusion into differential cohomology holds at the level of chains and is natural, we have

$$d(i(\phi)) = id(\phi) = i(Sq^{n-1}(\bar{a}_n)) = \hat{a}_n \cup \hat{a}_n. \quad (4.4)$$

Now the following matrix organizes the defining system given by equations (4.3) and (4.4):

$$\begin{pmatrix} 0 & \hat{a}_n & i(\phi) & \\ 0 & 0 & \hat{a}_n & \hat{A}_{(2n-1)} \\ 0 & 0 & 0 & \hat{b}_n \\ 0 & 0 & 0 & 0 \end{pmatrix},$$

and an element of the Massey product $\langle \hat{a}_n, \hat{a}_n, \hat{b}_n \rangle$ is given by the class

$$\hat{a}_n \cup \hat{A}_{(2n-1)} - i(\phi) \cup \hat{b}_n$$

We now consider the more interesting trivializations of String, Fivebrane [42] and Ninebrane structures [41]. In fact, what we will consider are slightly weaker versions, i.e. the vanishing of the p_i , $i = 1, 2, 3$, where p_i is the i th Pontrjagin class rather than the vanishing of the precise fractional classes. These differ from p_i -structures by the fact that we still require the lower Pontrjagin classes to vanish (see [41] for more discussion). We will then in turn consider differential refinements of these

structures, leading to Massey products representing *geometric* String, Fivebrane and Ninebrane structures, respectively (see Remark 9).

The following three examples are of the same type, but we separate for clarity. They involve sphere bundles with fiber S^{n-1} whose structure groups are $SO(n)$. Hence we can consider the corresponding Pontrjagin classes.

Example 14 (String structures). Now consider 3-sphere bundle $S^3 \rightarrow E \xrightarrow{\pi} M^4$ with nontrivial first Pontrjagin class $p_1(E) \neq 0$. Assuming that $H^2(M^4) \neq 0$, we take two elements $a_2, b_2 \in H^2(M^4)$ such that $\pi^*p_1(E) = a_2 \cup b_2$. Then we have a 2-connection on the S^3 bundle whose 3-form C_3 satisfies $dC_3 = \pi^*p_1(E) = a_2 \cup b_2$. As in Example 10 we can again assume that C_3 represents an integral cochain. Since $H^4(M^4)$ maps to 0 in $H^4(E)$, we also have the existence of a class D_3 such that $dD_3 = a_2 \cup a_2$ and we can form the Massey triple product $\langle a_2, a_2, b_2 \rangle \in H^5(E)$ which is represented by $D_3 \cup b_2 - a_2 \cup C_3$.

Example 15 (Fivebrane structures). Now consider 7-sphere bundle $S^7 \rightarrow E \xrightarrow{\pi} M^8$ with nontrivial second Pontrjagin class $p_2(E) \neq 0$. Assuming that $H^4(M^8) \neq 0$, we take two elements $a_4, b_4 \in H^4(M^8)$ such that $\pi^*p_2(E) = a_4 \cup b_4$. Then we have a 6-connection on the S^7 bundle whose 7-form C_7 satisfies $dC_7 = \pi^*p_2(E) = a_4 \cup b_4$. Choosing a singular cochain D_7 such that $dD_7 = a_4 \cup a_4$, we can again form the Massey product $\langle a_4, a_4, b_4 \rangle \in H^{11}(E)$ which is represented by $D_7 \cup b_4 - a_4 \cup C_7$.

Example 16 (Ninebrane structures). Now consider 11-sphere bundle $S^{11} \rightarrow E \xrightarrow{\pi} M^{12}$ with nontrivial third Pontrjagin class $p_3(E) \neq 0$. Assuming that $H^6(M^{12}) \neq 0$, we take two elements $a_6, b_6 \in H^6(M^{12})$ such that $\pi^*p_3(E) = a_6 \cup b_6$. Then we have a 10-connection on the S^{11} bundle whose 11-form C_{11} satisfies $dC_{11} = \pi^*p_3(E) = a_6 \cup b_6$. Choosing a singular cochain D_{11} such that $dD_{11} = a_6 \cup a_6$, we can again form the Massey product $\langle a_6, a_6, b_6 \rangle \in H^{17}(E)$ which is represented by $D_{11} \cup b_6 - a_6 \cup C_{11}$.

The above classical examples can be generalized and cast succinctly in the following. The proof is evident from the above examples.

Proposition 18 (p_n -structures). Consider sphere bundle $S^{4n-1} \rightarrow E \xrightarrow{\pi} M^{4n}$ with nontrivial n th Pontrjagin class $p_n(E) \neq 0$. Assuming that $H^{2n}(M^{4n}) \neq 0$, we take two elements $a_{2n}, b_{2n} \in H^{2n}(M^{4n})$ such that $\pi^*p_n(E) = a_{2n} \cup b_{2n}$. Then we have a $(4n-2)$ -connection on the S^{4n-1} bundle whose $(4n-1)$ -form C_{4n-1} satisfies $dC_{4n-1} = \pi^*p_n(E) = a_{2n} \cup b_{2n}$. Choosing a singular cochain D_{4n-1} such that $dD_{4n-1} = a_{2n} \cup a_{2n}$, we can again form the Massey product $\langle a_{2n}, a_{2n}, b_{2n} \rangle \in H^{6n-1}(E)$ which is represented by $D_{4n-1} \cup b_{2n} - a_{2n} \cup C_{4n-1}$.

We now give the differentially refined version of the proposition and provide the manifestations of the refined Massey products in each of the above examples. The proof follows directly from our earlier definitions and construction.

Proposition 19 (Differential p_n -structures). Consider the sphere bundle $S^{4n-1} \rightarrow E \xrightarrow{\pi} M^{4n}$ with classes $a_{2n}, b_{2n} \in H^{2n}(M^{4n})$. Let \hat{a}_{2n} and \hat{b}_{2n} be differential refinements such that $d\hat{C}_{4n-1} = \pi^*\hat{p}_n(E) = \hat{a}_{2n} \cup \hat{b}_{2n}$ and $d\hat{D}_{4n-1} = \hat{a}_{2n} \cup \hat{a}_{2n}$ so that the bundles $\hat{a}_{2n} \cup \hat{b}_{2n}$ and $\hat{a}_{2n} \cup \hat{a}_{2n}$ are trivializable with trivializations \hat{C}_{4n-1} and \hat{D}_{4n-1} . Then we can form the differential Massey product $\langle \hat{a}_{2n}, \hat{a}_{2n}, \hat{b}_{2n} \rangle \in \hat{H}^{6n-1}(E)$ with representative $\hat{D}_{4n-1} \cup \hat{b}_{2n} - \hat{a}_{2n} \cup \hat{C}_{4n-1}$.

Remark 9. Notice that from this proposition, we can construct Massey products from trivializations of differential String, Fivebrane and Ninebrane structures. In these cases, we can view the corresponding classes as representing geometric String, Fivebrane [43] and Ninebrane structures [41], respectively. Explicitly, these classes are

- (Geometric String) $\hat{D}_3 \cup \hat{b}_2 - \hat{a}_2 \cup \hat{C}_3 \in \langle \hat{a}_2, \hat{a}_2, \hat{b}_2 \rangle$.
- (Geometric Fivebrane) $\hat{D}_7 \cup \hat{b}_4 - \hat{a}_4 \cup \hat{C}_7 \in \langle \hat{a}_4, \hat{a}_4, \hat{b}_4 \rangle$.
- (Geometric Ninebrane) $\hat{D}_{11} \cup \hat{b}_6 - \hat{a}_6 \cup \hat{C}_{11} \in \langle \hat{a}_6, \hat{a}_6, \hat{b}_6 \rangle$.

4.2 Characteristic forms and anomaly cancellations

Consider a bundle P with curvature F on a manifold M . Let $c_i(P)$ be a characteristic class of degree i and let $c_i(F)$ be the corresponding characteristic form. Consider the conditions in cohomology $c_i(P) \cup c_j(P) = 0$ and $c_j(P) \cup c_k(P) = 0$. Then at the level of characteristic forms we have the trivializations via differential forms α and β of the indicated degrees

$$\begin{aligned} c_i(F) \wedge c_j(F) &= d\alpha_{(i+j-1)} , \\ c_j(F) \wedge c_k(F) &= d\beta_{(j+k-1)} . \end{aligned} \quad (4.5)$$

We build the composite differential form

$$\mu = c_i(F) \wedge \beta_{(j+k-1)} + (-1)^{i-1} \alpha_{(i+j-1)} \wedge c_k(F) \in \Omega^{i+j+k-1}(M) , \quad (4.6)$$

which is directly verified to be closed. This then allows us to form the Massey triple product of the corresponding cohomology classes

$$\langle c_i(P), c_j(P), c_k(P) \rangle \in H^{i+j+k-1}(M; \mathbb{Z}) . \quad (4.7)$$

Notice that we can consider conditions analogous to (4.5) in differential cohomology

$$\begin{aligned} \hat{c}_i(F) \cup \hat{c}_j(F) &= d\hat{\alpha}_{(i+j-1)} , \\ \hat{c}_j(F) \cup \hat{c}_k(F) &= d\hat{\beta}_{(j+k-1)} , \end{aligned} \quad (4.8)$$

requiring not only that the characteristic forms vanish, but that the corresponding bundles trivialize as bundles with connection. In this case, we can form the bundle (differential cochain)

$$\hat{\mu} = \hat{c}_i(F) \cup \hat{\beta}_{(j+k-1)} + (-1)^{i-1} \hat{\alpha}_{(i+j-1)} \cup \hat{c}_k(F) \in \text{Map}(M, \mathbb{B}^{i+j+k-2}U(1)_{\text{conn}}) , \quad (4.9)$$

which is an element in

$$\langle \hat{c}_i(P), \hat{c}_j(P), \hat{c}_k(P) \rangle \in \hat{H}^{i+j+k-1}(M; \mathbb{Z}) . \quad (4.10)$$

we summarize the above.

Proposition 20. Given a system (4.8) of trivializations of products of differential characteristic classes, we can build the stacky Massey product given by (4.9).

We now provide an application of this direct but fairly general observation. Consider a 10-dimensional manifold X^{10} with metric g on which there is a vector bundle with connection A . One can consider the setting in families, i.e. take a bundle E with fiber X^{10} and base a parameter space and then integrate over the fiber to get a class on the parameter space (see [22] for beautiful constructions). We will not do all this but simply just set up integral expressions which will suffice for our purposes. The Green-Schwarz anomaly polynomials are given as

$$\begin{aligned} I_4 &= p_1(g) - \text{ch}_2(A) , \\ I_8 &= -\text{ch}_4(A) + \frac{1}{48}p_1(g)\text{ch}_2(A) - \frac{1}{64}p_1(g)^2 + \frac{1}{48}p_2(g) . \end{aligned}$$

In [43] the first polynomial I_4 is interpreted as giving rise to a twisted String structure, and the indecomposable terms $p_2(g)$ and $\text{ch}_2(A)$ in I_8 are interpreted as giving rise (essentially) to a Fivebrane structure and its twist, respectively. Their trivializations H_3 and H_7 provide trivializations of String and Fivebrane structures, respectively. A question remained on how to interpret the decomposable terms in I_8 , namely $\frac{1}{48}p_1(g)\text{ch}_2(A)$ and $-\frac{1}{64}p_1(g)^2$. We provide one interpretation of the corresponding trivializations, which fits well within our context. Consider the situation when $[p_1 \cup \text{ch}_2] = 0 = [p_1 \cup p_1]$, i.e.

$$\text{ch}_2(A) \wedge p_1(g) = d\alpha_7(A, g) , \quad (4.11)$$

$$p_1(g) \wedge p_1(g) = d\beta_7(g) , \quad (4.12)$$

and build the differential form

$$\mu_{11} = \text{ch}_2(A) \wedge \beta_7(g) - \alpha_7(A, g) \wedge p_1(g) . \quad (4.13)$$

This form is closed by virtue of (4.11) and (4.12). Therefore, we can form the Massey triple product

$$\langle \text{ch}_2, p_1, p_1 \rangle \in H^{11}(E; \mathbb{Z}) . \quad (4.14)$$

As expected, the previous discussion refines to differential cohomology. Let X^{10} be as before. Since we are fixing a Riemannian metric on X^{10} and equipping the vector bundle with a connection A , it follows by uniqueness of characteristic forms [6],[45] that we have unique differential refinements

$$\begin{aligned} \hat{I}_4 &= \hat{p}_1(g) - \hat{\text{ch}}_2(A) , \\ \hat{I}_8 &= -\hat{\text{ch}}_4(A) + \frac{1}{48}\hat{p}_1(g)\hat{\text{ch}}_2(A) - \frac{1}{64}\hat{p}_1(g)^2 + \frac{1}{48}\hat{p}_2(g) . \end{aligned}$$

We now consider the situation when $[\hat{p}_1 \cup \hat{\text{ch}}_2] = 0 = [\hat{p}_1 \cup \hat{p}_1]$, i.e.

$$\hat{\text{ch}}_2(A) \wedge \hat{p}_1(g) = d\hat{\alpha}_7(A, g) , \quad (4.15)$$

$$\hat{p}_1(g) \wedge \hat{p}_1(g) = d\hat{\beta}_7(g) . \quad (4.16)$$

We then build the bundle

$$\hat{\mu}_{11} = \hat{\text{ch}}_2(A) \cup \hat{\beta}_7(g) - \hat{\alpha}_7(A, g) \cup \hat{p}_1(g) \in \text{Map}(X^{10}, \mathbb{B}^{10}U(1)_{\text{conn}}) \quad (4.17)$$

which is a representative of the Massey triple product

$$\langle \hat{\text{ch}}_2, \hat{p}_1, \hat{p}_1 \rangle \in \hat{H}^{11}(E; \mathbb{Z}) . \quad (4.18)$$

Therefore, we have the following

Proposition 21. *The mixed terms in the Green-Schwarz anomaly polynomials give rise to a stacky Massey product given by the top class (4.18).*

It is interesting to note the form of the connection on the bundle $\hat{\mu}$. Using the formula for the DB cup product, we see that the connection is

$$CS_3(A) \wedge CS_3(g) \wedge p_1(g) - \alpha_7(A, g) \cup p_1(g) , \quad (4.19)$$

which we will make use of below.

Fiber integration of Massey products and anomaly line bundles In [19] [20], a fiber integration map was defined by taking the usual fiber integration in cohomology, lifting to differential cohomology and then lifting to the internal hom in sheaves of positively graded chain complexes to produce a map

$$\int_{\Sigma^k} - : [N(C(\{U_i\}), \mathbb{Z}_{\mathcal{D}}^{\infty}[n]) \rightarrow \mathbb{Z}_{\mathcal{D}}^{\infty}[n-k] .$$

Here Σ^k is a paracompact manifold of dimension k and $C(\{U_i\})$ is the Čech nerve corresponding to a good open cover of Σ^k . The lifts are provided by the construction of Gomi and Terashima in [26]. Post-composing with the quasi-isomorphism provided by the exponential and applying the Dold-Kan functor gives a morphism of stacks in the form of holonomy

$$\text{hol}_{\Sigma^k} := \exp \left(2\pi i \int_{\Sigma_k} - \right) : [\Sigma^k, \mathbb{B}^n U(1)_{\text{conn}}] \rightarrow \mathbb{B}^{n-k} U(1)_{\text{conn}} . \quad (4.20)$$

Again in [19] [20], it was observed that the abelian Chern-Simons action functional can be described by post-composing the cup product morphism with this holonomy map. In particular, for a manifold Σ^{4k+3} , this composite induces an intersection pairing on differential cohomology

$$(\hat{x}, \hat{y}) \rightarrow \exp \left(2\pi i \int_{\Sigma^{4k+3}} \hat{x} \cup \hat{y} \right) . \quad (4.21)$$

For $k = 0$ and $\hat{y} = \hat{x}$, this pairing gives the usual Chern-Simons action. We now would like to describe how to lift this morphism to the Massey product (when defined). In fact, when the differential Massey product is defined, it is already a section

$$\langle \hat{x}, \hat{y}, \hat{z} \rangle \in [\Sigma^k, \mathbb{B}^{n_1+n_2+n_3+2} U(1)_{\text{conn}}] . \quad (4.22)$$

Hence, we can apply the fiber integration map to get the following.

Proposition 22. *The integration over the fiber of the differential Massey product (4.22) is the section*

$$e^{(2\pi i \int_{\Sigma^k} \langle \hat{x}, \hat{y}, \hat{z} \rangle)} \in \mathbb{B}^{n_1+n_2+n_3+2-k} U(1)_{\text{conn}} .$$

Remark 10. *Note the following.*

1. *The above construction can be generalized to higher Massey products, as we can fiber integrate any differential cohomology class of any degree, including those that are Massey products.*

2. The notation $e^{(2\pi i \int_{\Sigma^k} \langle \hat{x}, \hat{y}, \hat{z} \rangle)}$ is slightly abusive, since this map may not be well-defined on the entire Massey product (due to indeterminacy). What we really mean here is an element of the Massey product.

In particular, when \hat{x} , \hat{y} and \hat{z} come as characteristic forms, they are given by *morphisms* of stacks; e.g.

$$\hat{x} : [\Sigma^k, \mathbb{B}G_{\text{conn}}] \rightarrow [\Sigma^k, \mathbb{B}^{n_1}U(1)_{\text{conn}}] ,$$

which gives a natural assignment of differential cohomology classes as we vary the G -principal bundle with connection on Σ^k [19]. In this case, fiber integration gives the morphism of stacks

$$e^{(2\pi i \int_{\Sigma^k} \langle \hat{x}, \hat{y}, \hat{z} \rangle)} : [\Sigma^k, \mathbb{B}G_{\text{conn}}] \rightarrow \mathbb{B}^{n_1+n_2+n_3+2-k}U(1)_{\text{conn}} . \quad (4.23)$$

One interesting instance of this morphism comes from the previous example of Green-Schwarz anomaly polynomials. That is, we are interested in the triple product $\langle \hat{\text{ch}}_2, \hat{p}_1, \hat{p}_1 \rangle$. In this case, we get a morphism

$$e^{(2\pi i \int_{X^{10}} \langle \hat{\text{ch}}_2, \hat{p}_1, \hat{p}_1 \rangle)} : [X^{10}, \mathbb{B}G_{\text{conn}}] \rightarrow \mathbb{B}^{10-10}U(1)_{\text{conn}} = C^\infty(-, U(1)) , \quad (4.24)$$

from the moduli stack of bundles on X^{10} equipped with connection to smooth $U(1)$ -valued functions. It is useful to unwind this map at the level of connections. Indeed, we have at that level:

Proposition 23. *The morphism (4.23) is given by the exponential of*

$$\int_{X^{10}} \langle \hat{\text{ch}}_2, \hat{p}_1, \hat{p}_1 \rangle = \int_{X^{10}} CS_3(A) \wedge CS_3(g) \wedge p_1(g) - \alpha_7(A, g) \cup p_1(g) .$$

Remark 11. *Note the following.*

1. The exponential of the functional on the right, being built out of Chern-Simons forms, is indeed in $U(1)$.
2. As the structure of the functional in the proposition involves a product of two Chern-Simons forms, this suggests a formulation where X^{10} is viewed as a manifold of corners of codimension two, in the sense of the setting in [39] [40]. We hope to take up this point of view elsewhere.

4.3 Transfer of Massey products

We now relate the stacky Massey triple product to the triple Deligne-Beilinson cup product. To that end, we recall the following from [31] (the argument there was for specific dimensions but it extends evidently to any dimension). Consider Z^{n+1} as obtained from gluing two cobordisms together, i.e. Z^{n+1} is an orientable compact manifold and Y^n is a submanifold of codimension 1 such that $Z^{n+1} - Y^n$ has two connected components, each of which is a cobordism. Then from the Mayer-Vietoris sequence, there is a connecting (or transfer) map

$$T : H^k(Y^n) \rightarrow H^{k+1}(Z^{n+1}) . \quad (4.25)$$

Now let $a, b, c \in H^*(Z^{n+1})$ with restrictions $a', b', c' \in H^*(Y^n)$, and suppose further that the cup products vanish $a' \cup b' = b' \cup c' = 0 \in H^*(Y^n)$ so that the Massey product is defined. Then, by

considering the Poincaré dual chains, one has that the transfer of the Massey product gives the triple product [31]

$$T\langle a', b', c' \rangle = a \cup b \cup c \quad \text{mod indeterminacy} \quad (4.26)$$

where the Massey product is taken in $H^*(Y^n)$, the product in $H^*(Z^{n+1})$. The indeterminacy can be taken as $a \cup z + x \cup c$ where z, x are cocycles in the opposite connected components of $Z^{n+1} - Y^n$.

We now generalize this to our stacky setting of differential cohomology.

Proposition 24. *The connecting homomorphism for differential cohomology takes the form*

$$T : H^{k-1}(Y^n, U(1)) \rightarrow \hat{H}^{k+1}(Z^{n+1}),$$

and sends the differential Massey product to the triple DB cup product (modulo indeterminacy).

Proof. That the connecting homomorphism takes this form follows from the Mayer-Vietoris sequence for the sheaf hypercohomology of $\mathbb{Z}_D^\infty[n+1]$ [6]. Explicitly, the connecting map can be described as follow. Let U and V be an open cover of Z^{n+1} with $U \cap V \sim Y^n$ and let α be a flat Čech-Deligne cocycle on $U \cap V$. Now take a partition of unity $\{\rho_U, \rho_V\}$ subordinate to the cover $\{U, V\}$. Then the connecting map is given by

$$T(\alpha) = \begin{cases} -[D(\rho_U \alpha)] & \text{on } U \\ [D(\rho_V \alpha)] & \text{on } V. \end{cases}$$

In particular, if $\alpha \in \langle \hat{a}_1, \hat{a}_2, \hat{a}_3 \rangle$ is a representative of the Massey product, then α is of the form

$$\alpha = \hat{a}_1 \cup_{DB} \hat{x} + (-1)^k \hat{y} \cup_{DB} \hat{a}_3,$$

where $D\hat{x} = \hat{a}_2 \cup_{DB} \hat{a}_3$ and $D\hat{y} = \hat{a}_1 \cup_{DB} \hat{a}_2$ on $U \cap V$. Then it is straightforward to check directly from the explicit description of T that $T(\langle \hat{a}_1, \hat{a}_2, \hat{a}_3 \rangle) = \hat{a}_1 \cup_{DB} \hat{a}_2 \cup_{DB} \hat{a}_3$. \square

The Deligne-Beilinson triple cup product arises in the description of certain Chern-Simons type field theories in [19] [20]. The above result then can be applied to these theories to give the following, in the same notation as above.

Proposition 25. *The Massey triple product of three differential cohomology elements on Z^{n+1} transfers to a triple cup product Chern-Simons theory (in the sense of [19] [20]) on Y^n .*

4.4 Twisted cohomology and twisted Bianchi identities

We consider the Ramond-Ramond (RR) fields in type IIA string theory on a ten-dimensional manifold X^{10} with a B-field, whose curvature is a closed three-form H_3 . The RR fields of various degrees can be combined into the expression $\mathcal{F} = \sum_{i=0}^5 F_{2i}$, and satisfy the twisted Bianchi $d\mathcal{F}_n + H_3 \wedge \mathcal{F}_{n-2} = 0$. In components,

$$\begin{aligned} H_3 \wedge F_0 &= -dF_2, \\ H_3 \wedge F_2 &= -dF_4, \\ H_3 \wedge F_4 &= -dF_6, \\ H_3 \wedge F_6 &= -dF_8, \\ H_3 \wedge F_8 &= -dF_{10}, \end{aligned} \quad (4.27)$$

while $dF_0 = 0 = dF_{10}$. From these we will build expressions of degree ten.

1. Considering the first and fifth expressions in (4.27), we can set up the top differential form

$$\mu_1 = F_0 \wedge F_{10} + F_2 \wedge F_8 . \quad (4.28)$$

This is closed by dimension reasons, so that we can form the triple Massey product

$$\langle F_0, H_3, F_8 \rangle \in H^{10}(X^{10}; \mathbb{Z}) . \quad (4.29)$$

2. Considering the second and fourth expressions in (4.27), we build the top form

$$\mu_2 = F_2 \wedge F_8 + F_4 \wedge F_6 . \quad (4.30)$$

This is closed again by dimension reasons, and we can build the triple Massey product

$$\langle F_2, H_3, F_6 \rangle \in H^{10}(X^{10}; \mathbb{Z}) . \quad (4.31)$$

We now would like to refine the previous discussion to differential cohomology. Notice that since $dF_{2i} \neq 0$, we cannot simply put hats everywhere and expect the equations to hold at the level of ordinary differential cohomology. Consequently, there are two directions we can go. First, we could try to form Massey products in *twisted* differential cohomology, which is outside the scope of the present paper. Second, we can view the F_{2i} 's as *improved* gauge invariant field strengths corresponding to potentials C_{2i-1} with curvatures G_{2i} , which are not gauge invariant. We will expand on this latter point of view. To this end, we require that the potentials C_{2i-1} satisfy

$$dC_n + H_3 \wedge C_{n-2} = 0 \quad (4.32)$$

Notice that this equation implies that the improved field strengths F_{2i} vanish, by definition. Combining the potentials into the single potential $\mathcal{C} = \sum_{i=0}^3 C_{2i-1}$ we have, by assumption, the equations

$$\begin{aligned} H_3 \wedge C_1 &= -dC_3 , \\ H_3 \wedge C_3 &= -dC_5 , \\ H_3 \wedge C_5 &= -dC_7 . \end{aligned} \quad (4.33)$$

These equations can be viewed as conditions on the connections for differential refinements of the field strengths G_{2i} . Indeed, the full differentially refined equations read

$$\begin{aligned} \hat{H}_3 \cup \hat{G}_2 &= -d\hat{G}_4 , \\ \hat{H}_3 \cup \hat{G}_4 &= -d\hat{G}_6 , \\ \hat{H}_3 \cup \hat{G}_6 &= -d\hat{G}_8 . \end{aligned} \quad (4.34)$$

1. Considering the first and third expressions in (4.34), we can form the bundle

$$\hat{\mu}_1 = \hat{G}_2 \cup \hat{G}_8 + \hat{G}_4 \cup \hat{G}_6 , \quad (4.35)$$

with higher connection

$$C_1 \wedge G_8 + C_3 \wedge G_6 = C_1 \wedge H_3 \wedge C_5 + C_3 \wedge H_3 \wedge C_3 . \quad (4.36)$$

This bundle is an element in the stacky Massey triple product

$$\langle \hat{G}_2, \hat{H}_3, \hat{G}_6 \rangle \in \hat{H}^{10}(X^{10}; \mathbb{Z}) . \quad (4.37)$$

2. Considering instead the first and second expressions in (4.34), we form the higher bundle

$$\hat{\mu}_2 = \hat{G}_4 \cup \hat{G}_6 + \hat{G}_4 \cup \hat{G}_4 . \quad (4.38)$$

with higher connection

$$C_3 \wedge G_6 + C_5 \wedge G_4 = C_3 \wedge H_3 \wedge C_3 + C_5 \wedge H_3 \wedge C_1 . \quad (4.39)$$

This is an element in the stacky triple Massey product

$$\langle \hat{G}_4, \hat{H}_3, \hat{G}_4 \rangle \in \hat{H}^{10}(X^{10}; \mathbb{Z}) . \quad (4.40)$$

Proposition 26. *The system of twisted Bianchi identities for the differential RR fields leads to two higher bundles with connections (4.36) and (4.39) which are elements in the stacky Massey products in top degree (4.37) and (4.40), respectively.*

It would be interesting to investigate the implications of these expressions to string theory. For now we just observe that, essentially and up to signs, μ_1 and μ_2 are part of the couplings that arise in calculating the topological partition function of the RR fields (in the case when $H_3 = 0$) [12] [4]. While we do not pursue this here, we expect $\hat{\mu}_1$ and $\hat{\mu}_2$ to be relevant for the calculation of the partition function in the twisted differential case, $\hat{H}_3 \neq 0$, as well, extending the twisted topological case in [37] [36].

4.5 Quadruple Massey products

We now consider a setting inspired by type IIB string theory. Consider four fields $h_3^{(i)} \in H^3(X; \mathbb{Z})$, $i = 1, \dots, 4$, on a ten-dimensional manifold X^{10} , and consider analogs of three composite (Ramond-Ramond) fields $F_5^{(j)}$, $j = 1, 2, 3$, such that

$$\begin{aligned} h_3^{(1)} \wedge h_3^{(2)} &= -dF_5^{(3)} , \\ h_3^{(2)} \wedge h_3^{(3)} &= -dF_5^{(1)} , \\ h_3^{(3)} \wedge h_3^{(4)} &= -dF_5^{(2)} . \end{aligned}$$

Then there are further composite (analogs of Ramond-Ramond) fields $F_7^{(i)}$, $i = 1, \dots, 4$, such that

$$\begin{aligned} F_5^{(3)} \wedge h_3^{(3)} &= -dF_7^{(3)} , \\ h_3^{(1)} \wedge F_5^{(1)} &= -dF_7^{(1)} , \\ F_5^{(1)} \wedge h_3^{(4)} &= -dF_7^{(4)} , \\ h_3^{(2)} \wedge F_5^{(2)} &= -dF_7^{(2)} . \end{aligned}$$

Then we will end up (see below) having the Massey quadruple product as the integer

$$\langle h_3^{(1)}, h_3^{(2)}, h_3^{(3)}, h_3^{(4)} \rangle := -F_7^{(3)} \wedge h_3^{(4)} - F_7^{(1)} \wedge h_3^{(4)} - F_5^{(3)} \wedge F_5^{(2)} + h_3^{(1)} \wedge F_7^{(4)} + h_3^{(1)} \wedge F_7^{(2)} \in H^{10}(X^{10}; \mathbb{Z}) \cong \mathbb{Z} .$$

We now elaborate on the above. We first start with the triple Massey product in the current IIB string theory inspired context. Let $[h_3^{(i)}] \in H^3(X^{10})$ ($i = 1, 2, 3$) be non-zero cohomology classes such that $[h_3^{(1)}] \cup [h_3^{(2)}] = 0$ and $[h_3^{(2)}] \cup [h_3^{(3)}] = 0$. For the cocycle representatives $h_3^{(i)}$, write

$$h_3^{(1)} \cup h_3^{(2)} = dF_5^{(1)}, \quad (4.41)$$

$$h_3^{(2)} \cup h_3^{(3)} = dF_5^{(2)}. \quad (4.42)$$

Notice that from these two equations one gets immediately that $d(F_5^{(1)} \cup h_3^{(3)} + h_3^{(1)} \cup F_5^{(2)}) = 0$ by a straightforward application of the Leibnitz rule. This can then be used to define the triple Massey product as the subset of $H^8(X^{10})$ given by

$$\langle [h_3^{(1)}], [h_3^{(2)}], [h_3^{(3)}] \rangle = \left\{ \left[F_5^{(1)} \cup h_3^{(3)} + h_3^{(1)} \cup F_5^{(2)} \right] \right\}, \quad (4.43)$$

where h_3 and F_5 run over all possible choices above. The indeterminacy in the choice of the representative $w = h_3^{(1)} \cup F_5^{(2)} + F_5^{(1)} \cup h_3^{(3)}$ for the triple product lies in the ideal $([h_3^{(1)}], [h_3^{(2)}])$.

In order to connect with the Massey 4-fold product, it is good to rewrite the triple product in matrix form. The classes $[h_3^{(i)}]$ and the elements $F_5^{(i)}$ can be encoded in a matrix form

$$\begin{pmatrix} 0 & a_{11} & a_{22} & \\ & 0 & a_{23} & \\ & & 0 & a_{33} \\ & & & 0 \end{pmatrix} = \begin{pmatrix} 0 & h_3^{(1)} & F_5^{(1)} & \\ & 0 & h_3^{(2)} & F_5^{(2)} \\ & & 0 & h_3^{(3)} \\ & & & 0 \end{pmatrix}. \quad (4.44)$$

The defining properties of the Massey triple product can be expressed in a matrix multiplication as

$$\begin{aligned} d \begin{pmatrix} 0 & h_3^{(1)} & F_5^{(1)} & \\ & 0 & h_3^{(2)} & F_5^{(2)} \\ & & 0 & h_3^{(3)} \\ & & & 0 \end{pmatrix} &= \begin{pmatrix} 0 & 0 & h_3^{(1)} h_3^{(2)} & \\ & 0 & 0 & h_3^{(2)} h_3^{(3)} \\ & & 0 & 0 \\ & & & 0 \end{pmatrix}, \\ \begin{pmatrix} 0 & h_3^{(1)} & F_5^{(1)} & \\ & 0 & h_3^{(2)} & F_5^{(2)} \\ & & 0 & h_3^{(3)} \\ & & & 0 \end{pmatrix}^2 &= \begin{pmatrix} 0 & 0 & h_3^{(1)} h_3^{(2)} & h_3^{(1)} F_5^{(2)} + F_5^{(1)} h_3^{(3)} \\ & 0 & 0 & h_3^{(2)} h_3^{(3)} \\ & & 0 & 0 \\ & & & 0 \end{pmatrix}. \end{aligned} \quad (4.45)$$

Now in order to go one step further to the quadruple (or 4-fold) product, we need to satisfy certain conditions on the triple product, in analogy to saying that higher obstructions arise only once the lower ones vanish. So in our case, we first need to assume that we can complete our set by adding two more elements, a fourth $h_3^{(4)}$ and a third $F_5^{(3)}$, such that

$$dF_5^{(3)} = h_3^{(3)} \cup h_3^{(4)}. \quad (4.46)$$

Besides the above representative w , we then have a second representative for the triple product and is given by $z = h_3^{(2)} \cup F_5^{(3)} + F_5^{(2)} \cup h_3^{(4)}$, namely representing $\langle [h_3^{(2)}], [h_3^{(3)}], [h_3^{(4)}] \rangle$. The condition

to be able to define the quadruple product is that *both* triple products vanish *simultaneously*, i.e. that both cohomology representatives w and z can be chosen as coboundaries, which we write as $w = dF_7^{(1)}$ and $z = dF_7^{(2)}$.

We are now ready to define the 4-fold or quadruple Massey product. In analogy to the triple product, we start with the equations (4.41) (4.42) and (4.46), and then write the two cocycles of degree eight

$$dF_7^{(1)} = h_3^{(1)} \cup F_5^{(2)} + F_5^{(1)} \cup h_3^{(3)} , \quad (4.47)$$

$$dF_7^{(2)} = h_3^{(2)} \cup F_5^{(3)} + F_5^{(2)} \cup h_3^{(4)} , \quad (4.48)$$

from which we get a cocycle

$$x = h_3^{(1)} \cup F_7^{(2)} + F_5^{(1)} \cup F_5^{(2)} + F_7^{(1)} \cup h_3^{(4)} \quad (4.49)$$

of degree ten. Again, define the quadruple Massey product $\langle [h_3^{(1)}], [h_3^{(2)}], [h_3^{(3)}], [h_3^{(4)}] \rangle$ as a collection of all cohomology classes $[x] \in H^{10}(X^{10})$ that we can obtain by the above procedure.

The indeterminacy is best presented as the *matrix* triple product of certain elements, namely of $(h_3^{(1)}, H^5(X^{10}))$, $\begin{pmatrix} h_3^{(2)} & H^5(X^{10}) \\ 0 & h_3^{(3)} \end{pmatrix}$, and $\begin{pmatrix} H^5(X^{10}) \\ h_3^{(4)} \end{pmatrix}$.

We now generalize this construction to differential cohomology. To produce the desired products, we again view the $F_5^{(i)}$ and $F_7^{(i)}$ as improved, gauge invariant field strengths and denote the corresponding potentials as $C_4^{(i)}$ and $C_6^{(i)}$, with curvatures $G_5^{(i)}$ and $G_7^{(i)}$. We now lift everything to the level of differential cohomology, which yields the equations

$$\begin{aligned} \hat{h}_3^{(1)} \cup \hat{h}_3^{(2)} &= -d\hat{G}_5^{(3)} , \\ \hat{h}_3^{(2)} \cup \hat{h}_3^{(3)} &= -d\hat{G}_5^{(1)} , \\ \hat{h}_3^{(3)} \cup \hat{h}_3^{(4)} &= -d\hat{G}_5^{(2)} , \end{aligned} \quad (4.50)$$

and

$$\begin{aligned} \hat{G}_5^{(3)} \cup \hat{h}_3^{(3)} &= -d\hat{G}_7^{(3)} , \\ \hat{h}_3^{(1)} \cup \hat{G}_5^{(1)} &= -d\hat{G}_7^{(1)} , \\ \hat{G}_5^{(1)} \cup \hat{h}_3^{(4)} &= -d\hat{G}_7^{(4)} , \\ \hat{h}_3^{(2)} \cup \hat{G}_5^{(2)} &= -d\hat{G}_7^{(2)} . \end{aligned} \quad (4.51)$$

The connection on the higher bundle is calculated as follows:

$$\begin{aligned} &-C_6^{(3)} \wedge h_3^{(4)} - C_6^{(1)} \wedge h_3^{(4)} - C_4^{(3)} \wedge G_5^{(2)} + b_2^{(1)} \cup G_7^{(4)} + b_2^{(1)} \cup G_7^{(2)} = \\ &= -C_6^{(3)} \wedge h_3^{(4)} - C_6^{(1)} \wedge h_3^{(4)} - C_4^{(3)} \wedge b_2^{(3)} \wedge h_3^{(4)} + b_2^{(1)} \wedge C_4^{(1)} \wedge h_3^{(4)} + b_2^{(1)} \wedge b_2^{(2)} \wedge b_2^{(3)} \wedge h_3^{(4)} . \end{aligned}$$

Here $b_2^{(i)}$ denotes a local potentials for the forms $h_3^{(i)}$. Therefore, we have the following description as phase or holonomy.

Proposition 27. *The system (4.50) (4.51) leads to the stacky Massey quadruple product*

$$\langle \hat{h}_3^{(1)}, \hat{h}_3^{(2)}, \hat{h}_3^{(3)}, \hat{h}_3^{(4)} \rangle := -\hat{G}_7^{(3)} \cup \hat{h}_3^{(4)} - \hat{G}_7^{(1)} \cup \hat{h}_3^{(4)} - \hat{G}_5^{(3)} \cup \hat{G}_5^{(2)} + \hat{h}_3^{(1)} \cup \hat{G}_7^{(4)} + \hat{h}_3^{(1)} \cup \hat{G}_7^{(2)} \in \hat{H}^{10}(X^{10}; \mathbb{Z}),$$

viewed as a higher bundle with connection given by

$$-C_6^{(3)} \wedge h_3^{(4)} - C_6^{(1)} \wedge h_3^{(4)} - C_4^{(3)} \wedge b_2^{(3)} \wedge h_3^{(4)} + b_2^{(1)} \wedge C_4^{(1)} \wedge h_3^{(4)} + b_2^{(1)} \wedge b_2^{(2)} \wedge b_2^{(3)} \wedge h_3^{(4)}.$$

The discussion using matrix Massey products carries over to differential cohomology in a similar fashion.

Acknowledgement

The authors would like to thank Domenico Fiorenza and Urs Schreiber for very useful discussions and comments. We would also like to thank Chris Kapulkin for useful comments on the first version of the manuscript, and the referee for useful suggestions.

References

- [1] I. K. Babenko and I. A. Taimanov, *Massey products in symplectic manifolds*, Sb. Math. **191** (2000) 1107, [arXiv:math.SG/9911132].
- [2] C. Bär and C. Becker, *Differential Characters*, Lecture Notes in Mathematics 2112, Springer, 2014.
- [3] A. Beilinson, *Notes on absolute Hodge cohomology*, Applications of algebraic K-theory to algebraic geometry and number theory, Part I, II, Contemp. Math. 55, Amer. Math. Soc., Providence, RI, 1986.
- [4] D. M. Belov and G. W. Moore, *Type II actions from 11-dimensional Chern-Simons theories*, [arXiv:hep-th/0611020].
- [5] J. Brylinski, *Loop spaces, characteristic classes and geometric quantization*, Progress in Mathematics, Vol. **107**, Birkhäuser Boston, 2008.
- [6] U. Bunke, *Differential cohomology*, [arXiv:math.AT/1208.3961].
- [7] U. Bunke and Th. Schick, *Uniqueness of smooth extensions of generalized cohomology theories*, J. Topol. **3** (2010), no. 1, 110–156, [arXiv:0901.4423].
- [8] U. Bunke, *Differential function spectra, the differential Becker-Gottlieb transfer and application to differential algebraic K theory*, [arXiv:math.At/1306.0247].
- [9] J. Cheeger and J. Simons, *Differential characters and geometric invariants*, Lecture Notes in Math. 1167 (1985), 55–90, Springer-Verlag, New York. 1985.
- [10] K. T. Chen, *Connections, holonomy and path space homology*, in Differential geometry (Proc. Sympos. Pure Math., Vol. XXVII, Part 1, Stanford Univ., Stanford, Calif., 1973), pp. 39–52, Amer. Math. Soc., Providence, R. I., 1975.

- [11] K. T. Chen, *Free subalgebras of loop space homology and Massey products*, Topology **11**, 3 (1972), 237–243.
- [12] E. Diaconescu, G. Moore and E. Witten, *E_8 gauge theory, and a derivation of K -theory from M -theory*, Adv. Theor. Math. Phys. **6** (2003) 1031–1134, [arXiv:hep-th/0005090].
- [13] P. Deligne, *Théorie de Hodge II*, IHES Pub. Math. **40** (1971), 5–57.
- [14] P. Deligne and D. S. Freed, *Classical field theory*, in Quantum fields and strings: a course for mathematicians, vol. 1 (Princeton, NJ, 1996/1997), 137–225, Amer. Math. Soc., Providence, RI, 1999.
- [15] P. Deligne, P. Griffiths, J. Morgan, and D. Sullivan, *The real homotopy theory of Kähler manifolds*, Inventiones Math. **29** (1975), 245–254.
- [16] C. Deninger, *Higher order operations in Deligne cohomology*, Inventiones Math. **122** (1995), 289–315.
- [17] D. Dugger, *Combinatorial model categories have presentations*, Adv. Math. **164** (2001), no. 1, 177–201, [arXiv:math/0007068] [math.AT].
- [18] H. Esnault and E. Viehweg, *Deligne-Beilinson cohomology*, in Beilinson’s conjectures on special values of L-functions, 43–91, Perspect. Math., 4, Academic Press, Boston, MA, 1988.
- [19] D. Fiorenza, H. Sati, and U. Schreiber, *Extended higher cup-product Chern-Simons theory*, J. Geom. Phys. **74** (2013), 130–163, [arXiv:1207.5449] [hep-th].
- [20] D. Fiorenza, H. Sati, and U. Schreiber, *A Higher stacky perspective on Chern-Simons theory*, Mathematical Aspects of Quantum Field Theories (Damien Calaque and Thomas Strobl eds.), Springer, Berlin (2015), [arXiv:1301.2580] [hep-th].
- [21] D. Fiorenza, U. Schreiber, and J. Stasheff, *Cech cocycles for differential characteristic classes – An infinity-Lie theoretic construction*, Adv. Theor. Math. Phys. **16** (2012), 149–250, [arXiv:1011.4735] [math.AT].
- [22] D. S. Freed, *Dirac charge quantization and generalized differential cohomology*, Surveys in differential geometry, 129–194, Surv. Differ. Geom., VII, Int. Press, Somerville, MA, 2000, [arXiv:hep-th/0011220].
- [23] P. Gajer, *Geometry of Deligne cohomology*, Invent. Math. **127** (1997), no. 1, 155–207.
- [24] P.G. Goerss and J.F. Jardine, *Simplicial Homotopy Theory*, Birkhäuser, Basel, 2009.
- [25] K. Gomi, *Differential characters and the Steenrod squares*, Adv. Stud. Pure Math. **52** (2008) 297–308, [arXiv:math/0411043] [math.AT].
- [26] K. Gomi and Y. Terashima, *A fiber integration formula for the smooth Deligne cohomology*, Internat. Math. Res. Notices **13** (2000), 699–708.
- [27] D. Grady, *Massey products in differential cohomology*, PhD thesis, University of Pittsburgh, 2015.

- [28] P. Griffiths and J. Morgan, *Rational Homotopy Theory and Differential Forms*, Birkhäuser, NY, 2013.
- [29] M. J. Hopkins, I. M. Singer, *Quadratic functions in geometry, topology, and M-theory*, J. Diff. Geom. **70** (2005), 329–452, [arXiv:math/0211216].
- [30] D. Kraines, *Massey higher products*, Trans. Amer. Math. Soc. **124** (1966), 431–449.
- [31] I. Kriz and H. Sati, *Type II string theory and modularity*, J. High Energy Phys. **0508** (2005) 038, [arXiv:hep-th/0501060].
- [32] J. Lurie, *Higher topos theory*, Princeton University Press, 2009.
- [33] W. S. Massey, *Some higher order cohomology operations*, International symposium on algebraic topology, Universidad Nacional Autónoma de Mexico and UNESCO, Mexico City, 1958, 145–154.
- [34] W. S. Massey, *Higher order linking numbers*, Conf. Algebraic Topology, Univ. of Illinois at Chicago Circle, 1968, pp. 174–205.
- [35] J. P. May, *Matric Massey products*, J. Algebra **12** (1969), 533–568.
- [36] V. Mathai and H. Sati, *Some relations between twisted K-theory and E_8 gauge theory*, J. High Energy Phys. **03** (2004) 016, [arXiv:hep-th/0312033].
- [37] G. Moore and N. Saulina, *T-duality, and the K-theoretic partition function of type IIA superstring theory*, Nucl. Phys. **B 670** (2003), 27–89, [arXiv:hep-th/0206092].
- [38] M. Morishita and Y. Terashima, *Geometry of polysymbols*, Math. Res. Lett. **15** (2008), no. 1, 95–115.
- [39] H. Sati, *Corners in M-theory*, J. Phys. **A44** (2011), 255402, [arXiv:1101.2793].
- [40] H. Sati, *M-theory with framed corners and tertiary index invariants*, SIGMA **10** (2014), 024, [arXiv:1203.4179].
- [41] H. Sati, *Ninebrane structures*, Int. J. Geom. Methods in Modern Phys. **12** (2015) 1550041, [arXiv:1405.7686] [hep-th].
- [42] H. Sati, U. Schreiber, and J. Stasheff, *Fivebrane structures*, Rev. Math. Phys. **21** (2009) 1–44, [0805.0564] [math.AT].
- [43] H. Sati, U. Schreiber, and J. Stasheff, *Differential twisted String- and Fivebrane structures*, Commun. Mathematical Phys. **315** (2012), 169–213, [0910.4001] [math.AT].
- [44] A. Schwarzhaupt, *Massey products in Deligne-Beilinson cohomology*, PhD thesis, Essen University, 2002.
- [45] J. Simons and D. Sullivan, *An axiomatic characterization of ordinary differential cohomology*, J. Topology **1** (1) (2008), 45–56, [arXiv:math/0701077].
- [46] J. Stasheff, *H-spaces from a Homotopy Point of View*, Springer LNM 161, 1970.

- [47] U. Schreiber, *Differential cohomology in a cohesive infinity-topos*, [[arXiv:1310.7930](#)] [math-ph].
- [48] T. Wenger, *Massey products in Deligne-cohomology*, PhD thesis, Münster University, 2000.